

USE AREA	PESTS	APPLICATION RATE	NOTES
ROSES & FLOWERS: (1)	Aphids, Mites, Leaf-hoppers, Thrips, Luonymus Scale Crawlers, Flea Beetles, Dip-terous Leaf Miners, Mealybugs, Whiteflies	4 tbsps. (2 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray	Spray underside of leaves. Wet plants to the dripping point
WREDS & SHRUBS:	Aphids, Mites, Scale Crawlers, Dipterous Leaf Miners, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Leafhoppers, Mealybugs, Obscure Root Weevils, Omnivorous Leaf Tiers	4 tbsps. (2 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray.	Wet leaves and branches to the dripping point. Try to penetrate dense foliage.
	Webworms, (Juniper, Mimosa), Whiteflies	4 tbsps. (2 fl. oz.) in 3 gals. of water. Apply as a thorough foliar spray.	Mimosa Webworms: Apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.
	Apple and Thorn Ske-letonzers, Hemlock, Caperes, Oak Loopers, Oblique-banded Leaf Rollers, Pear Stugs, Tent Caterpillars, Webworms.	6 oz. in 3 gals. of water. Apply as a thorough foliar spray.	

(1) Do not use on certain Ferns, Hibiscus, Poinsettia, Papaya, Pilea, Gardenia and Orchids, since injury may occur.

Fruit Spray: Mix 16 oz. in 24 gallons of water to control Codling Moths, Mites, Aphids, Leafrollers and Scale Crawlers on Apples, Peaches, Plums, Grapes, Cherries and Strawberries. Start spraying after blooming and repeat at 10-14 day intervals if needed. Do not spray within 21 days of picking.

Vegetable Spray: Mix 4 tbsps. (2 fl. oz.) in 3 gallons of water (16 oz. in 24 gallons). Three gallons of diluted spray will cover 1000 sq. ft. of Tomatoes, Carrots, Potatoes, Beans, Cucumbers, Onions, Cantaloupes, Peppers and Cabbage to control Bean Beetles, Aphids, Leaf Miners, Imported Cabbageworms, and Cucumber Beetles. Make applications at 7-10 day intervals when insects appear. Do not spray within 14 days of picking. Tomatoes may be sprayed 1 day before picking. Potatoes may be sprayed 35 days before harvest. Wash home garden vegetables before using.

Nuisance Pests in Outside Areas:

Ants*, Boxelder Bugs, Brown Dog Ticks*, Chiggers*, Clover Mites*, Cockroaches, Crickets*, Earwigs*, Fleas*, Flies*, Millipedes*, Sawbugs (Dillbugs)*, Spiders, Springtails (Collembola)*, 1. 16 fl. oz. in 3 gals. of water.

Thoroughly spray window and door screens, sills, foundations, porch and patio walls and ceilings, entrance walks, garbage cans, tree trunks and into cracks and other places where insects can hide.

Do not treat plants or animals.

*Spray a 5 ft. band of soil around the house next to the foundation of the house to a height of 2-3 feet.

†See lawn recommendations for other control measures for these pests.

Fire Ants:

To aid in the control of fire ants in lawns and other recreation areas, apply 2 fl. oz. (4 tbsp.) in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration reducing product effectiveness.

Note: For best results, apply in cool weather of 65-80°F or in early morning or late evening hours. Treat new mounds as they appear.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSAL: 20, 35 & 55 gal. drums

a) Reseal container and offer for reconditioning or b) triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place.

SMALLER CONTAINERS: Do not reuse empty container. Wrap container and put in trash collection.

GENERAL: Consult Federal, State or local disposal authorities for approved alternative procedures.

BEST AVAILABLE COPY

397

3 3

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals. Avoid storage near food or feed products. Do not contaminate ornamental fishponds.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Do not induce vomiting.

If Inhaled: Remove victim to fresh air and apply artificial respiration if indicated.

If on Skin: Wash promptly with soap and water. Rinse thoroughly.

If in Eyes: Rinse eyes with plenty of water and call a physician if irritation persists.

Note to Physician: If symptoms of cholinesterase inhibition occur, Atropine Sulfate is antidotal and may be administered in conjunction with 2-PAM.

ENVIRONMENTAL HAZARDS



This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL HAZARDS

Do not use: pour, spill or store near heat or open flame

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

NET CONTENTS: _____



BEST AVAILABLE COPY

BATCH _____



Printed in U.S.A.