

4816-604

P11-17

101

CODE 722.35

# Pyrenone<sup>®</sup> INDUSTRIAL E.C. 60-6

EPA Reg. No. 4816-604 EPA Est. No. 279-NY-1

AN INSECTICIDE SPRAY CONCENTRATE FOR DILUTION WITH WATER OR OIL

FOR USE IN FOOD PROCESSING AND STORAGE, HOSPITALS, HOTELS, RESTAURANTS, OTHER INSTITUTIONS AND IN RESIDENCES.

Provides Rapid Kill Without Objectible Residues.

Authorized for use in all departments of official establishments operation under the meat, poultry, shell egg grading and egg products inspection programs.

The insecticide components (piperonyl butoxide and pyrethrins) are classified as food additives and when this product is used as directed, allowable tolerance levels will not be exceeded.

ACTIVE INGREDIENTS	
Pyrethrins .....	6.00%
*Piperonyl Butoxide, Technical .....	50.00%
**INERT INGREDIENTS .....	34.00%
100.00%	

Equivalent to 48-(butylcarbityl) (6-pronyl-miscronyl) ether and 127 related compounds.  
\*Contains Petroleum Distillate  
PYRENGU<sup>®</sup> - Registered Trademark of Fairfield American Corporation.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
See side panel for additional precautionary statements.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**BEST AVAILABLE COPY**

When diluting with water, prepare only the amount needed for immediate use. Holding a water dilution more than 24 hours is not recommended. Agitate occasionally. For oil dilutions, a deodorized petroleum oil is recommended.

For maximum effectiveness, a combination of localized surface application and space treatment is recommended.

AS A SURFACE SPRAY: For initial clean up of severe insect infestations, dilute at a rate of 1 part concentrate in 29 parts water or oil. This dilution may not be used in edible product areas of federally inspected meat and poultry plants. In these plants and for normal infestations, dilute 1 part concentrate in 59 parts water or oil.

To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, mealworms, grain mites and cadellies:  
Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles, spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

AS A SPACE SPRAY: Dilute at a rate of 1 part concentrate to 119 parts water or oil. Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths direct spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Ventilate and sweep up dead insects before reoccupying.

SPRAY: For rapid kill of exposed or accessible stages of the insects named on this label, dilute 1 part concentrate to 11 to 59 parts water or oil. Apply using conventional mechanical or compressed air equipment (not thermal) following directions for space spraying. Apply the above dilutions at ratio from 1 to 10 ounces per 1000 cubic feet of space.

When used in meat and poultry processing plants, a dilution of no less than 1 to 11 (5.0%) is acceptable and it must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns).

SEE REVERSE SIDE

**FAIRFIELD AMERICAN CORPORATION**  
3937 Salt Road  
Medina, New York

2831 LEM

