

For Use in Bakeries, Bottling Plants, Canneries,  
Flour Mills, Food Processing Plants, Grain  
Elevators, Hospitals, Hotels, Motels, Poultry  
Houses, Restaurants, and Outdoors for Mos-  
quito Control and on Growing Crops.

May Be Applied As A Wet Spray In Food Plants  
While In Operation.

Pyrethrins.....	3.00%
*Piperonyl Butoxide, Technical....	30.00%
Petroleum Distillate.....	12.00%
INERT INGREDIENTS:.....	55.00%
	100.00%

**Keep Out Of Reach Of Children**

See side panel for additional precautions.

EPA Reg. No. 4816-605  
EPA Est. No. 279-NY-1

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To prepare dilutions the concentrate should first be stirred or agitate well. Add the required amount of concentrate to water and blend thoroughly. This will produce a stable micro-emulsion.

The insecticidal components of this concentrate, pyrethrins and piperonyl butoxide, are classified as food additives, and when used as directed allowable tolerance levels will not be exceeded.

To control exposed stages of houseflies, including fruit flies, mosquitoes, gnats, and small flying moths dilute 1 part concentrate in 59 parts water (2.1 ounces per gallon). Close doors and windows and shut off ventilating systems. Apply as a fog or fine mist, directing the spray toward the upper portions of the room and toward windows and other light sources which attract these insects.

Thoroughly fill the area with mist using 1 to 2 ounces per 1000 cubic feet. Keep area closed for 10 minutes or more. Vacate after treatment and ventilate before reoccupying. Repeat treatment as needed.

For initial clean out of infestations of cockroaches, ants, spiders, silverfish, crickets, boxelder bugs, scorpions, and clover mites: Dilute 1 part concentrate in 14 parts water (8.5 fluid ounces per gallon). Apply as a coarse droplet surface spray using approximately 1 gallon to 750 square feet. Spray thoroughly all surfaces where these insects are usually found including walls, baseboards, moldings, doors, cabinets, shelves, sinks and pipes, with special attention to cracks and crevices, niches, dark corners, drains and similar hiding places. Treat ant trails, nests and runs. When possible, spray insects directly.

Repeat treatment using a dilution of 1 plus 29 (4.3 fluid ounces per gallon) at the same rate of application after an interval of one month or when reinfestation occurs.

IN MILLS, ELEVATORS AND GRANARIES: For rapid initial kill or exposed or accessible stages of granary weevil, rice weevil, confused flour beetles, rust red flour beetle, cadelle, sawtoothed grain beetle, cheese mites and grain mites. Dilute at a rate of 1 part concentrate to 29 parts water (4.3 fluid ounces per gallon). Clean out any waste grain, dust, dirt or debris and destroy before spraying. Apply as a coarse droplet spray using approximately 1 gallon to 750 square feet of surface.

Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this use.

In mills, elevators, and granaries, thoroughly spray floors, walls and other surfaces of bins, storage and handling areas with particular attention to cracks, crevices, and similar protected locations. Treat unloading, handling and processing areas, and apply the insecticide to conveying, processing and handling equipment such as the inside of reel conveyors, purifier conveyors, sifters, roll housings and hoppers, pick up conveyors, etc. Spray thoroughly around the base of heavy machinery and equipment.

Thoroughly treat walls, floors, shelves and other exposed surfaces, and lightly spray each layer of cartons or containers as stacked.

Sack surfaces of bagged grain can be sprayed lightly until slightly moist.

For protection in transit, treat truck beds, box cars, and ships' holds before loading.

Using a dilution of 1 part concentrate in 29 parts water (4.3 fluid ounces per gallon), apply at a rate of 1 gallon to 750 square feet of surface with particular attention to cracks, crevices, and similar hiding places.

ACCEPTED  
NOV 19 1980  
F. Edith Harrison,  
and Raymond A. R.  
for the patient  
No. 7716-635

[illegible]

# FOR

# FOR



**FAIRFIELD  
AMERICAN  
CORPORATION**

23 CORPORATION  
3932 Salt Road  
Medina, New York 14103

BEST DOCUMENT AVAILABLE

11 4816-605

PM 17

103 Code 720,000

# PYRENONE<sup>®</sup>

## MULTI-PURPOSE AQUEOUS 30-3

BEST AVAILABLE COPY

### A HIGHLY VERSATILE PYRENONE INSECTICIDE CONCENTRATE

For Use in Bakeries, Bottling Plants, Canneries, Flour Mills, Food Processing Plants, Grain Elevators, Hospitals, Hotels, Motels, Poultry Houses, Restaurants, and Outdoors for Mosquito Control and on Growing Crops.

Provides Rapid Kill of Insects Without Objectionable Residues.

May Be Applied As A Wet Spray In Food Plants While In Operation.

#### ACTIVE INGREDIENTS

Pyrethrins.....	3.00%
*Piperonyl Butoxide, Technical.....	30.00%
*INERT INGREDIENTS.....	67.00%
	100.00%

\*Equivalent to 26% (butylcarbitol)(6-propyl-piperonyl) ether and to 6% related compounds.

\*\*Contains Petroleum Distillate

Keep Out Of Reach Of Children

**CAUTION**

See side panel for additional precautions.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To prepare dilutions the concentrate should first be stirred or agitate well. Add the required amount of concentrate to water and blend thoroughly. This will produce a stable micro-emulsion.

The insecticidal components of this concentrate, pyrethrins and piperonyl butoxide, are classified as food additives, and when used as directed allowable tolerance levels will not be exceeded.

#### INDOOR APPLICATION:

To control exposed stages of houseflies, including fruit flies, mosquitoes, gnats, and small flying moths dilute 1 part concentrate in 59 parts water (2.1 ounces per gallon). Close doors and windows and shut off ventilating systems. Apply as a fog or fine mist, directing the spray toward the upper portions of the room and toward windows and other light sources which attract these insects.

Thoroughly fill the area with mist using 1 to 2 ounces per 1000 cubic feet. Keep area closed for 10 minutes or more. Vacate after treatment and ventilate before reoccupying. Repeat treatment as needed.

For initial clean out of infestations of cockroaches, ants, spiders, silverfish, crickets, boxelder bugs, scorpions, and clover mites: Dilute 1 part concentrate in 14 parts water (8.5 fluid ounces per gallon). Apply as a coarse droplet surface spray using approximately 1 gallon to 750 square feet. Spray thoroughly all surfaces where these insects are usually found including walls, baseboards, moldings, doors, cabinets, shelves, sinks and pipes, with special attention to cracks and crevices, niches, dark corners, drains and similar hiding places. Treat ant trails, nests and runs. When possible, spray insects directly.

Repeat treatment using a dilution of 1 part concentrate in 29 (4.3 fluid ounces per gallon) at the same rate of application after an interval of one month or when reinfestation occurs.

**IN MILLS, ELEVATORS AND GRANARIES:** For rapid initial kill of exposed or accessible stages of granary weevil, rice weevil, confused flour beetle, rust red flour beetle, cadelle, sawtoothed grain beetle, cheese mites and grain mites. Dilute at a rate of 1 part concentrate to 29 parts water (4.3 fluid ounces per gallon). Clean out any waste grain, dust, dirt or debris and destroy before spraying. Apply as a coarse droplet spray using approximately 1 gallon to 750 square feet of surface.

Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this use.

In mills, elevators, and granaries, thoroughly spray floors, walls and other surfaces of bins, storage and handling areas with particular attention to cracks, crevices, and similar protected locations. Treat unloading, handling and processing areas, and apply the insecticide to conveying, processing and handling equipment such as the inside of reel conveyors, purifier conveyors, sifters, roll housings and hoppers, pick up conveyors, etc. Spray thoroughly around the base of heavy machinery and equipment.

Thoroughly treat walls, floors, shelves and other exposed surfaces, and lightly spray each layer of cartons or containers as stacked.

Sack surfaces of bagged grain can be sprayed lightly until slightly moist.

For protection in transit, treat truck beds, box cars, and ships' holds before loading.

Using a dilution of 1 part concentrate in 29 parts water (4.3 fluid ounces per gallon), apply at a rate of 1 gallon to 750 square feet of surface with particular attention to cracks, crevices, and similar hiding places.

PYRENONE - Registered Trademark of Fairfield American Corporation

EPA Reg. No. 4816-605  
EPA Est. No. 279-NY-1

See Reverse Side

ACCEPTED

FAIR  
AME  
CORPO  
3932 Sall  
Medina, N