

pm 10 4816-633

page 1 of 1

CODE 72470

PYRENONE®

FLEA & TICK SPRAY

EPA Reg. No. 4816-633AA EPA Est. No. 279-NY-1

* FOR USE ON CATS AND DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS

* AN AQUEOUS, STABLE, MICRO EMULSION
* ALCOHOL FREE—PREVENTS DRY SKIN

ACTIVE INGREDIENTS:

Pyrethrins.....	0.06%
*Piperonyl Butoxide, Technical.....	0.60%
INERT INGREDIENTS:.....	99.34%
	100.00%

*Equivalent to min. 0.48% (butylcarbityl) (6-propylpiperonyl) ether and 0.12% related compounds.

PYRENONE - Registered Trademark of Fairfield American Corporation

KEEP OUT OF REACH OF CHILDREN
5 MAR 1993
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4816-633

NET CONTENTS

ROUSSEL UGLAF
 Roussel Uclaf Corporation
 95 Chestnut Ridge Road
 Montvale, New Jersey 07645



72470 103090E

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USE

ON PETS

Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and under body. While spraying, fluff the hair so that spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes, face, or genitals. Repeat as necessary. Do not spray on nursing females or pets under six weeks of age.

ON LIVESTOCK

To protect cattle, horses and hogs from *Horn Flies*, *House Flies*, *Mosquitoes* and *Gnats* apply a light mist sufficient to wet the surface of the hair. To control *Stable Flies*, *Horse Flies* and *Deer Flies*, apply at a rate of 2 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control *Blood Sucking Lice*, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

To control *Poultry Lice*, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of *Bedbugs* and *Mites* in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the *Bedbugs* and *Mites* hide.

ON PREMISES

To control *Fleas* and *Brown Dog Ticks* on premises: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding

should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a cool dry place inaccessible to children and pets and away from heat and direct sunlight. Protect from freezing.

DISPOSAL: Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center. Induce vomiting by touching back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES OR ON SKIN: Flush affected areas with plenty of water. Contact a physician if irritation persists.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid inhalation of vapors. Wash contaminated skin promptly with soap and water after use. Avoid spraying food or foodstuffs. Remove birds and cover fish aquariums before spraying.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

CODE 7247