





AIMM

Association for Information and Image Management

1100 Wayne Avenue Suite 1100  
Silver Spring Maryland 20910



301/587-8202




Centimeter



Inches



MANUFACTURED TO AIMM STANDARDS  
BY APPLIED IMAGE, INC.



# BUTACIDE 8 EC

EPA Reg. No. 4816-642AA

EPA Est. No. 279-NY-1

- \* **Tank-Mix Insecticide Synergist**
- \* **Contains 8 pounds of Piperonyl Butoxide per gallon**
- \* **Improves control of Resistant Insects**
- \* **For use in Resistance Management Programs**

## ACTIVE INGREDIENT:

Piperonyl Butoxide Technical

91.0%

## INERT INGREDIENTS:

9.0%

100.0%

\*Equivalent to 72.8% (butylcarbitol)(6-propylpiperonyl) ether and 18.2% related compounds

BUTACIDE - Registered trademark of Fairfield American Corporation

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED:** Remove affected person to fresh air. Apply artificial respiration if indicated.

**IF ON SKIN:** Wash with plenty of water and soap.

**IF IN EYES:** Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

See back of label for directions

### NET CONTENTS



FAIRFIELD  
AMERICAN  
CORPORATION

2010-11-17 10:00:00 AM

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### USE ON AGRICULTURAL CROPS:

To be tank mixed with agricultural insecticides, as recommended by local Agricultural Extension Specialists, County Extension Agents, Crop Consultants or Pest Control Advisors. To provide increased control of resistant insects, tank mix Butacide 8 EC at a rate of up to 1 pound of Piperonyl Butoxide per acre (see dilution chart below) with the appropriate amount of companion insecticide and sufficient diluent to ensure uniform coverage and apply per the Directions for Use on the companion insecticide label. Butacide 8 EC may be applied by ground or aerial application equipment.

### DILUTION CHART

Pounds PB Per Acre	Fluid Ounces Per Acre	Acres per Gallon
0.125	2	64
0.25	4	32
0.5	8	16
1.0	16	8

Applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

### USE THROUGH IRRIGATION SYSTEMS (CHEMIGATION)

Refer to supplemental labeling entitled "DIRECTIONS FOR APPLICATION THROUGH IRRIGATION SYSTEMS" (available through your Fairfield American distributor) for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed and the product to be tank mixed with is labeled for chemigation use.

### USE ON GREENHOUSE VEGETABLE, FLOWER, AND FOLIAGE PLANTS:

To be tank mixed with other insecticides to improve the control of the insects on the crops named on their labels. Observing the directions on the insecticide label, tank mix at a rate of up to 1 pound of Piperonyl Butoxide per 100 gallons of water (to provide 1 pound of Piperonyl Butoxide use 16 fluid ounces (1 pint) of Butacide 8 EC).

Plant safety is an important consideration when using insecticides in a greenhouse. However, it is not possible to evaluate the phytotoxicity of Butacide 8 EC in combination with a number of insecticides towards a multitude of plant varieties in different growth stages under varying environmental conditions. Before making widespread applications of Butacide 8 EC, treat a limited number of plants with the tank mix and observe for phytotoxicity over a 10-day period.

Applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

### USE ON ORNAMENTAL FLOWER, AND FOLIAGE PLANTS:

To be tank mixed with other insecticides on the crops named on their labels. Observing the directions on the insecticide label, tank mix at a rate of up to 1 pound of Piperonyl Butoxide per 100 gallons of water (to provide 1 pound of Piperonyl Butoxide use 16 fluid ounces (1 pint) of Butacide 8 EC).

Applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

### USE IN STRUCTURES

IN COMBINATION WITH OTHER INSECTICIDES, but not limited to, Baygon, Cyfluthrin, Diazinon, Dursban, Ficam, etc.

To provide increased control of insects in structures, tank mix 0.75 to 3.0 fluid ounces of Butacide 8 EC with the appropriate amount of companion insecticide and apply per the Directions for Use on the companion insecticide label.

Except in Federally inspected processing operations may be used as a surface spray with care and precautions given above.

In USDA inspected facilities, Butacide 8 EC with the appropriate amount of companion insecticide per finished gallon of water.

See back of label for directions

See back of label for directions

Percent Piperonyl Butoxide

0.5  
1.0  
2.0

Applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

Baygon, Commodore, Conquer, Cynoff, Demon, Dursban, Ficam, Pynd, Salfotin and Tempo are Registered Trademarks of Miles Inc., ICI Americas Inc., Paragon Professional Pest Control Products, FMC Corporation, ICI Americas Inc., Dow Chemicals, Nor-Am Chemical Company, Terminix International, Inc., Zococon Corporation and Miles Inc., respectively.

## STORAGE AND DISPOSAL

### CONTAINERS LARGER THAN ONE GALLON:

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE AND SPILL PROCEDURES:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

### CONTAINERS ONE GALLON AND SMALLER:

**STORAGE:** Store in original container in a cool, dry place inaccessible to children and pets.

**DISPOSAL:** Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

Concentrated material causes temporary eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray. Wash thoroughly with soap and water after handling.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, ponds, streams, tidal marshes or estuaries. Do not contaminate water by loading of equipment or disposal of wastes.

## WARRANTY LIMITATIONS AND DISCLOSURE

**READ CAREFULLY:** Fairfield American Corporation warrants only that this product conforms to the chemical description on its label and is reasonably fit for the purposes stated on its label only when used as stated in the directions and under normal use conditions. Conditions beyond Fairfield American Corporation's control, such as weather conditions, the combination with or presence of other materials, the manner in which the product is used or applied, or other unforeseeable factors, may result in

unintended consequences, such as crop injury, or ineffectiveness. No warranty extends to use of the product not in accordance with directions or use under abnormal or unforeseeable conditions, or conditions outside Fairfield American Corporation's control, and Buyer assumes the risk of all such uses.

FAIRFIELD AMERICAN CORPORATION DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. Buyer's sole remedy for breach of this warranty shall be limited to refund of the purchase price. Fairfield American Corporation shall not be liable for any consequential, special or indirect damages, such as lost profits or values, which result from the use or handling of this product.

**ACCEPTED**  
**05 FEB 1992**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
is amended, for the pesticide  
registered under  
EPA Reg. No. 4816-642