

Pyrenone[®] PESTIGAS[®]

EPA Reg. No. 100-100000, EPA Est. No.

This unique aerosol application system employs LIQUID CARBON DIOXIDE, a solvent and propellant.

No chemicals to mix, measure or pour.

No containers to dispose of.

These cylinders are returned to the manufacturer for refilling.

Active Ingredients

Pyrethrin	0.5%
*Piperonyl Butoxide	1.0%
Petroleum Distillate	98.5%
Inert Ingredients	0.5%

*Equivalent to 0.1% of piperonyl butoxide (piperonyl butoxide) other than 82-00-00 compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautions.

Pyrenone[®] - Registered trademark of Fairfield American Corporation.
PESTIGAS[®] - Registered trademark of Fairfield American Corporation.

Use indoors and outdoors for application as a contact or surface spray.

INDOOR USE AREAS

Aircraft	Hotel
Bakeries	Hotel
Beverage Plants	Hotel
Bottling Plants	Hotel
Breweries	Hotel
Buses	Hotel
Cafes	Hotel
Canneries	Hotel
Cattle Barns	Hotel
Condominiums	Hotel
Dairies	Hotel
Dried Food Products	Hotel
In Storage or Processing	Hotel
Drug Store	Hotel
Flour Mills	Hotel
Food Processing Plant	Hotel
Garage	Hotel
Grain Elevator	Hotel
Granary	Hotel
Greenhouse	Hotel
Home	Hotel
Hotel	Hotel
Hospital	Hotel
Hotel	Hotel
Industrial Installations	Hotel

OUTDOOR USE AREAS

Cattle Barns	Hotel
Cattle	Hotel
Dried Food Products	Hotel
Drives or Treatments	Hotel
Feedlots	Hotel
Grain Elevators	Hotel

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Pyrenone Pesticide should be used only with the appropriate high pressure pump system through an approved automatic spray system.

Two separate nozzle systems are available for use. The first is a high pressure nozzle which delivers a fine spray of the active ingredients at a charge rate of 1.0 g per second. The second is a low pressure nozzle which delivers a coarse spray of the active ingredients at a charge rate of 10.0 g per second. The use of the low pressure nozzle will allow for a more effective insect control.

An automatic timing device can also be coupled with a high delivery rate nozzle (10.0 g/sec). This timer can be set to activate at predetermined times to allow the nozzle to operate for the given amount of time for the area and insect pests to be treated (usually 1/2 hour on ft.).

The LDRC with the extended nozzle provides for use as a contact or surface spray as well as crack and crevice or void treatment.

To prepare your Pyrenone Pesticide cylinder for application follow directions supplied with each high pressure gun and hose. Open the cylinder valve. The cylinder is now ready for use. Upon completion of treatment release cylinder valve and relieve pressure from gun and hose.

Use of this product in food processing or food service establishments (bakeries, breweries, dairies, etc.) for the control of insects, rodents, etc. Industrial Installations (e.g., grain elevators, silos, etc.) should be treated during periods when the plants or facilities are not in operation. Hands should be removed or covered during treatment. All surfaces should be covered during treatment or thoroughly cleaned before reusing.



INSTRUCTIONS

SPACE AND CONTACT SPRAY APPLICATION

To control accessible areas, use the following:

Angoumois Grain Moth	1.0 g/sec
Ants	1.0 g/sec
Boxelder Bugs	1.0 g/sec
Caddisflies	1.0 g/sec
Cheese Mites	1.0 g/sec
Cheese Skippers	1.0 g/sec
Cigarette Beetles	1.0 g/sec
Clover Mites	1.0 g/sec
Corn Ear Moth	1.0 g/sec
Confused Flour Beetle	1.0 g/sec
Crickets	1.0 g/sec
Dark Mealworms	1.0 g/sec
Drugstore Beetles	1.0 g/sec
Earwigs	1.0 g/sec
Firebrats	1.0 g/sec
Fleas	1.0 g/sec
Flies	1.0 g/sec
Fruit Flies	1.0 g/sec
Grain Mites	1.0 g/sec
Granary Weevils	1.0 g/sec

SEE REVERSE SIDE



FAIRFIELD AMERICAN

Close doors and windows and keep off ventilating system. Direct the spray into wall space and crevices, behind all equipment, and under obstructions or dark harborage areas to provide maximum penetration and impingement of the insects. Then apply with space spray, directing the spray toward the ceiling and upper corners of the area and behind all obstructions.

Using the LDK, apply at a rate of 4 seconds per 10,000 cubic feet of space.

Using the HDR, apply at a rate of 4 seconds per 50,000 cubic feet of space.

Keep area closed for at least one hour. Ventilate the treated area and ventilate entire room before reentry. Repeat treatment if necessary.

TREATMENT OF STORED FOOD AREAS

Food in multi-walled bags or in other containers stored in warehouses, storage rooms, mills, grain elevators and granaries.

To control accessible adult stages of

Caddis	Grain Weevils
Confused Flour Beetles	Cigarette Beetles
Rice Weevils	Grain Mites
Sawtoothed Grain Beetle	Grain Weevils
Yellow Mealworms	Dark Mealworms
Cheese Mites	Grain Mites
	As Flour Beetles

Infested stored products should be destroyed, fumigated or treated with other effective means using an approved product intended for this purpose. As noted above, otherwise prevent Pestigas from directly spraying into cracks and crevices, corners, under pallets and around stacks or packages from settling as fine insects as possible.

Using the LDK, apply at a rate of 4 seconds per 100 square feet of floor space.

Using the HDR, apply at a rate of 4 seconds per 100 square feet of floor space.

Leave the treated area closed for one hour. Do not reenter the treated area and ventilate before reentry. Repeat the application if necessary.

ANIMAL QUARTERS Use in cattle barns, horse barns, poultry houses, swine pens, kennels, etc.

As a space spray, it is effective against Flying Moths, Mosquitoes, etc.

Using the LDK, apply at the rate of 4 seconds per 10,000 cubic feet of space above the animals.

Using the HDR, apply at the rate of 4 seconds per 50,000 cubic feet of space above the animals.

Direct spray towards the upper portions of loose falling feed with mist. Keep area closed for at least 1 hour before ventilating the area and ventilate feed room before reentry. Avoid exposure of feed to flies.

Do not treat or cause exposure to children or puppies under 1 or weeks of age. Repeat treatment if reinfestation occurs.

IN MUSHROOM PRODUCTION AND PACKING

To control Mushroom Flies and House Flies

In Growing Houses

Using the LDK, apply at the rate of 4 seconds per 10,000 cubic feet of space.

Using the HDR, apply at the rate of 4 seconds per 50,000 cubic feet of space.

Guns should be discharged 8 to 10 feet away from the beds and directed slightly downward over them. Circulation fans should be kept in operation, but for best results, exhaustors should be turned off.

In Packing and Canning Plants

Using the LDK, apply as above at the rate of 4 seconds per 10,000 cubic feet of space.

Using the HDR, apply as above at the rate of 4 seconds per 50,000 cubic feet of space.

SURFACE SPRAY APPLICATION

To control accessible stages of

Ants	Flies
Boxelder Bugs	Millipedes
Clover Mites	Millipedes
Cryptine Beetles	Grain Weevils
Earwigs	Grain Weevils
Firebrats	Grain Weevils
Flies	Grain Weevils

Using the LDK with the extension nozzle, apply at the rate of 4 seconds per 100 square feet of floor space. Spray walls, floors, and other surfaces of bins and storage areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Treat unloading, handling, and storage areas and inside conveying, processing and packing equipment. Spray thoroughly the tops of heavy machinery and equipment where insects settle directly when in use.

FOR IN-TRANSIT PROTECTION Use in trucks, truck beds, boxcars, etc. before loading. Then apply to the walls and floors and spray each layer of material in containers lightly as stacked.

TREATMENT OF STORED GRAIN

To control accessible adult stages of

Indian Meal Moths, Angoumois Grain Moths, Mediterranean Flour Moths

Apply to the surface of bins and grain, spray walls, floors and other surfaces of bins and storage, and handling areas. The grain being especially carefully treated in cracks, crevices and similar hiding places. Avoid exposure of grain to flies.

towards the ceiling and upper corners of the area and behind any obstructions.

Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, treat up within with a rake before spraying, and make a second application in two weeks.

Using the LDK, apply at a rate of 4 seconds per 10,000 cubic feet of space.

Using the HDR, apply at a rate of 4 seconds per 50,000 cubic feet of space.

TREATMENT OF FLYING INSECT RESTING PLACES

To aid in the control of:

Mosquitoes	Gnats
Small Flying Moths	Wasps

Treat walls, ceilings, moldings, screens, door frames, window frames, beams, etc. and similar resting places with the above directions as indicated for surface spray application.

Carpet Beetles: Spray directly on beetles, baseboards, floors, shelves and carpets. Treat both sides of rugs, carpets and pads and all exposed and accessible surfaces of overhead furniture. Use the LDK in combination with the extension nozzle and apply at a rate of 4 seconds per 100 square feet of floor space. Repeat treatment as necessary.

Bedbugs: Using the LDK in combination with the extension nozzle at a rate of 4 seconds per 100 square feet of floor space. Spray mattresses, particularly around tufts and seams. Turn bed apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

Flies: Using the LDK in combination with the extension nozzle at a rate of 4 seconds per 100 square feet of floor space. Spray sleeping quarters, bedding, floor and floor covering, where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

1. Treat Animals. To protect pet and dairy cattle from attacks of stable flies, horn flies, deer flies, house flies, barn flies, mosquitoes, the adult stages of blood-sucking flies and gnats, spray using the LDK in combination with the extension nozzle at a rate of 4 seconds per 100 square feet or about 1 lb. of material. Be very careful to spray back, withers and legs thoroughly. To protect from tick flies, spray the face and head, but not the ears and the eyes. Repeat treatment as necessary when flies are troublesome.

OUTDOOR USE

FOR TEMPORARY REDUCTION OF

House flies	Mosquitoes
Gnats	Stable flies

For use in drive-in theaters, drive-in restaurants, golf courses, urban areas, parks, playgrounds and other recreational areas. Direct the spray into tall grass, and around lawns, where these pests may hover or rest. Apply while the air is still and avoid wetting foliage. Application should be made prior to use. Repeat as necessary.

In the treatment of corrals, feedlots, swimming pools and zoos, cover any exposed water, draining fountains and animal feed before application. Fill the area with mist and direct the spray into tall grass, shrubbery and around lawns where the above pests may hover or rest. Apply while the air is still and avoid wetting foliage. In zoos, avoid exposure of reptiles to the product. Repeat as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: This cylinder must be checked periodically. A defective cylinder should not be used. A cylinder that has parts missing, is plainly worn, or is distorted, should be replaced immediately. Store in a cool, dry area.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSITION: This cylinder must not be disposed of in any manner. Cylinders must be returned to the distributor for credit of cylinder deposit or replacement. All cylinders, valves and caps remain the property of the distributor. Federal regulations prohibit the refilling of cylinders without written consent of the distributor.

SEE NEXT PAGE

**PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid inhalation of spray mist. Avoid contact with skin and eyes. Avoid contamination of feed and feedstuffs. Vacate premises after treatment and ventilate before reoccupying. Remove pets, birds and cover fish aquariums before spraying.

When using this product as a space spray in food processing plants, foods should be removed or covered during treatment. Do not apply as a space spray while food processing is underway. Except in federally inspected meat and poultry plants, when applied as a surface spray with care and in accordance with the directions and precautions given above food processing operations may continue. Thoroughly wash all food processing surfaces before re-use. All surfaces in processing areas must be washed and rinsed in potable water after spraying.

When using in animal quarters, do not apply directly to food, water or food supplements. Wash teats of dairy animals before milking.

STATEMENT OF PRACTICAL TREATMENT

- If in Eyes:** Flush eyes with plenty of water. Contact a physician if irritation persists.
- If Inhaled:** Remove affected person to fresh air. Apply respiration if indicated.
- If on Skin:** Remove contaminated clothing and wash affected areas with soap and water.
- If Swallowed:** Contact a physician or poison control center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes or estuaries. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Strip and at least may be killed at application rates recommended on this label. Do not apply where there are important resources.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Keep the container cool and open flame. Do not puncture or incinerate container. Keep safe to temperatures above 120°F. May cause burning.

Never use near oil tanks or oil storage. Handling of this product not to attract fire. Follow with directions given hereafter.

NET CONTENTS _____

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460		EPA REGISTRATION NO.	DATE OF ISSUANCE 10/13
NOTICE OF PESTICIDE: ☐ REGISTRATION ☐ REREISTRATION <i>Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended.</i>		TERM OF ISSUANCE	
		NAME OF PESTICIDE PRODUCT	
NAME AND ADDRESS OF REGISTRANT (include ZIP code) 16/18			
NOTE: Changes in labeling, formula, differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.			
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act. A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith. Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. (CAD)			
☐ ATTACHMENT IF APPLICABLE			
SIGNATURE OF APPROVING OFFICIAL			DATE