

Reg # 4816-666

PM-17

181

FSN 6840 INSECTICIDE, PYRETHRUM, SPACE SPRAY, SYNERGIZED PYRETHRUM.

Net Contents _____ Gallon(s)
Designed For Use By The Military

An Effective Semi-Concentrated PYRENONE[®]
Insecticide

For Application in Specialized Mechanical,
Thermal, or Compressed Air Fogging
Equipment.

Use Against House Flies, Mosquitoes,
Gnats, Midges, Wasps and Small Flying
Moths.

For Use in Bakeries, Food Plants, Res-
taurants, Hotels, Motels, Hospitals,
Schools, Office Buildings, Food Processing
or Handling Plants and Warehouses.

ACTIVE INGREDIENTS

Pyrethrins	0.4%
*Piperonyl Butoxide, Technical	1.6%
**INERT INGREDIENTS	98.0%
	100.0%

*Equivalent to min. 1.28% (butylcarbityl) (6-propylpiperonyl)
ether and 0.32% related compounds.

**Contains deodorized kerosene

PYRENONE - Registered trademark of
Fairfield American Corporation

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Contact a physician or
Poison Control Center. Do
not induce vomiting unless
directed by a physician.
This product contains a
high amount of petroleum
distillate. Vomiting may
cause aspiration pneumonia.

If Inhaled: Remove affected person to
fresh air. Apply artificial
respiration if indicated.

If in Eyes or on Skin: Flush with plenty
of water. Contact a physi-
cian if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not
apply directly to water. Do not contamin-
ate water by cleaning of equipment or dis-
posal of wastes. Apply this product only
as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour or spill near heat or
open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use
this product in a manner inconsistent with
its labeling.

This ready-to-use concentrate may be dis-
persed through good mechanical, thermal or
compressed air fogging or spraying equip-
ment. If used in edible product areas of
federally inspected meat, poultry, shell
egg grading and egg product establishments,
it must be dispersed as an aerosol spray
(no droplets over 50 microns in diameter
and 80% less than 30 microns).

To control house flies, mosquitoes, gnats,
midges and small flying moths, close room
and spray into the air. Adjust the nozzle
of the sprayer to give the finest mist
possible. Spray at a rate of 3/4 to 1
fluid ounce per 1000 cubic feet or until a
moderate mist can be seen in all parts of
the room. It is not necessary to spray the
air heavily. Keep the room closed for 10
minutes.

To control wasps using the above described
application, spray late in the evening
when insects are at rest.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by
storage or disposal.

PESTICIDE DISPOSAL:
Pesticide, spray mixture or rinsing water
that cannot be used according to label
instructions must be disposed of according
to applicable Federal, State or local
procedures.

CONTAINER DISPOSAL:
Triple rinse (or equivalent) then offer
for recycling or reconditioning, or punc-
ture and dispose of in a sanitary land-
fill, or by other approved State and
local authorities.

A container size of 30 gallons or more may
be resealed and offered for reconditioning.

Buyer assumes all risks of use, storage or
handling of this material not in strict
accordance with directions given herewith.

ACCEPTED
with conditions
in EPA Reg. No. 4816

OCT 1992

Under the
Federal
as well as
registered with Reg. No.
4816-666

EPA Reg. No. 4816- EPA Est. No. 279-NY 1

LOT NUMBER

Printed in U.S.A.

BEST AVAILABLE COPY

**KEEP OUT OF REACH OF CHILDREN
CAUTION**


PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

arm. If swallowed or absorbed through
skin. Avoid prolonged or repeated breath-
ing of vapors or spray mist. Avoid con-
tact with skin and eyes. Avoid contamina-
tion of feed and foodstuffs. In commercial
food handling facilities, cover or remove
all food and food processing equipment
prior to application. Do not apply while
food processing is underway. After space
spraying in bakeries and food processing
plants, wash all equipment, benches,
mixers, etc., where exposed food will be
handled, with potable water. All food
processing surfaces and utensils should be
washed during treatment, or thoroughly
washed before use.

Remove pets, birds, and cover fish aquariums
before spraying.



**FAIRFIELD
AMERICAN
CORPORATION**
3912 Salt Road
Medley, N.Y. 11761