



# Pyrenone<sup>®</sup> PRESSURIZED

EPA Reg. No. 4816-

A Water Based PYRENONE<sup>®</sup> Insecticide

Industrial Strength - Low Dosage Required

Use in Schools, Hospitals, Institutions, Hotels, Airports, Supermarkets, Restaurants, Greenhouses, Motels, Drugstores, Food Plants, Homes, Apartments, Theaters, Trucks, Camps, and Storage Buildings.

### ACTIVE INGREDIENTS

Pyrethrin	0.002
Pyrethrin Hydroxide, Technical	1.000
Petroleum Distillate	5.000
INERT INGREDIENTS	93.998
	100.000

Equivalent to min. 1.27 (0.01% active) (0.01% active) (0.01% active) other and 0.01% related compounds.

PYRENONE - Registered trademark of Fairfield American Corporation

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See side panel for additional precautions

NET CONTENTS \_\_\_\_\_

Printed in U.S.A.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE.**

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

Cockroaches, Earwiggs, Fleas, Waterbugs, Millipedes, Firebrats, Beetles, Millipedes, Spiders, Centipedes and Box Bugs: Spray into hiding places such as cracks and crevices, behind links, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray stem over possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Flies, Mosquitoes, Gnats, Midges, House Flies, and Flying Squirrels (Millers): Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in room 1 hour after treatment. Repeat treatment as necessary.

Bedbugs: Spray mattress, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, walling and frames. Repeat treatment as necessary.

For all flies, use Stray Bag: Take on legs, etc. and premises. Apply directly to animals along back, under forelegs and on other areas, except on animal's face. Do not contact skin and insure good coverage. Hold nozzle about one foot away while spraying. Apply until animal hair is thoroughly saturated. Do not spray directly in eyes or mouth of animal. Repeat within 24 hours if first treatment is necessary. To reduce reinfection thoroughly spray the animal's bedding and other resting areas until the sur-

face is slightly moist. Also spray the entire inside surface of dog houses and kennels, thoroughly spraying floors, cracks, crevices and bedding. Hit as many insects directly as possible. Repeat as necessary.

Indian Meal Moths and Angoumois Grain Moths: Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application as necessary.

Commoned Flour Beetles, Sawtoothed Grain Weevils, Angoumois Moths, House Weevils, and Flour Weevils: and Grain Mites: Spray into cracks and crevices, processing and handling equipment, and on cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 15 minutes, to 1 hour after treatment. Repeat application as necessary.

## DISPOSAL

Wrap container and put in trash collection. Do not incinerate.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

Avoid breathing of vapor or contact with skin or eyes. Vacate premises after treatment and ventilate before reoccupying.

Avoid contamination of feed and foodstuffs. When using this product in food processing plants, food should be removed or covered during treatment. All food processing surfaces should be cleaned during treatment and thoroughly cleaned and rinsed with potable water before using. When using this product in these areas, apply only when the facility is not in operation.

Remove pets, birds and cover fish aquarium before spraying.

PHYSICAL OR CHEMICAL HAZARDS: contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 100°F may cause bursting. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

4816-612



FAIRFIELD AMERICAN CORPORATION

BEST DOCUMENT AVAILABLE

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing plants, wash with potable water all equipment, benches, shelving, etc., where exposed food will be handled. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

*Joseph M. ... 26 October 1982*  
Franklin D. R. Gee  
Product Manager (17)  
Insecticide-Rodenticide Branch  
Registration Division (28-767) (2/203)

Enclosures