


PM 03

4816-687

6/27/86

186

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	EPA REGISTRATION NO. 4816-687 TERM OF ISSUANCE Until Reregistered NAME OF PESTICIDE PRODUCT Turbocide Pest Control System with DDVP	DATE OF ISSUANCE JUN 27 1986
NAME AND ADDRESS OF REGISTRANT (Include ZIP code) <div style="text-align: right; margin-right: 100px;">1611 173871 681</div> Fairfield American Corporation 8th & Harrison Streets Frenchtown, NJ 08825		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.		
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.		
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:		
1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.		
2. Make the labeling changes listed below before you release the product for shipment:		
a. Add the phrase "EPA Registration No. 4816-687." b. The total of item 13a on the Confidential Statement of Formula is 1000.1. Change 890.4 to 890.3. c. Delete "unique aerosol." Words or phrases implying that a product possesses unique characteristics are not acceptable. Change to read "This spray application system, etc." d. Change the percent of Related Compounds from 0.71 to 0.70. e. Move signal word DANGER to below the child hazard statement. f. Add skull and crossbones and the word POISON in red in close proximity to the signal word.		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL 	DATE 6/27/86	

- g. Place Note to Physician on front panel following STATEMENT of PRACTICAL TREATMENT.
- h. Add "SEE ADDITIONAL PRECAUTIONARY STATEMENTS on side panel" following Note to Physician.
- i. Make "in consistent" one word in the misuse statement.
- j. Change "Turbocide with DDVP" to "Turbocide Pest Control System with DDVP" or to "this product" wherever it appears in the Directions for Use.
- k. Restore PRECAUTIONARY STATEMENTS as on the draft label submitted February 4, 1985. See also paragraphs 7 and 8 of our letter dated November 13, 1985.
- l. Describe in the Precautionary Statements the preparation of treatment of enclosed places, use of approved respirator, and posting of signs.
- m. Very few of the pests listed under "And the Following Greenhouse or Horticultural Pests" are pests of greenhouse or horticulture. The group may be headed "And various stages of the following pests." Note that "mealybugs" is one word.
- n. Delete all references to "System II" and "Fixed Installation System." When these systems are ready to be marketed, submit brochures for our review. Along with the brochures you may request an amendment action requesting addition of these methods of application.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

TURBOCIDE PEST CONTROL SYSTEM WITH DDVP

EPA REG. NO. 4816-

EPA EST. NO. 38719-NC-1

For Sale only to Authorized Distributors and Approved Applicators.

This is a High Pressure Insect Spray

This unique aerosol application system employs liquid carbon dioxide as both solvent and propellant. There are no chemicals to mix, measure or pour and no container disposal. These cylinders are to be returned to the manufacturer for refilling.

NET CONTENTS:

ACTIVE INGREDIENTS:

*Dichlorvos [2,2-Dichlorovinyl dimethyl	phosphatel.....	9.30%
Related Compounds.....	0.71%	

INERT INGREDIENTS:

Liquid Carbon Dioxide.....	90.00%
----------------------------	--------

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

*Equivalent to 10.0% DDVP Insecticide

TURBOCIDE - is a Trademark of Fairfield American Corporation.

D A N G E R

KEEP OUT OF REACH OF CHILDREN

POISON

See Note to Physician and Additional Precautions on Side Panel

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Move victim to fresh air. Apply artificial respiration if indicated. Call a physician immediately.

IF ON SKIN: Remove contaminated clothing and wash immediately with soap and water. Call a physician immediately.

IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner in consistent with its labeling.

TurboCide with DDVP should be applied only with the appropriate high pressure gun and hose or through an approved spraying system.

SYSTEM I

The spray nozzle system provides for economical and accurate control of insect pests. The high pressure gun operates at a discharge rate of 2.0 g/sec (0.2 g of DDVP). The high pressure gun with the extended nozzle provides for use as a contact spray as well as a void treatment.

SYSTEM II

An automatic timing device can also be coupled with an approved spraying system. This timer can be set to be activated at pre-determined times to allow the nozzle to operate for the given amount of time for the area and insect pests to be treated. Refer to the System II brochure for specific operation.

FIXED INSTALLATION SYSTEM

Refer to the Fixed Installation brochure for specific operation.

To prepare your TurboCide with DDVP cylinder for application, follow directions supplied with each high pressure gun and hose. Open the cylinder valve. The cylinder is now ready for use. Upon completion of treatment, reclose cylinder valve and relieve pressure in gun and hose.

TurboCide with DDVP is a specially formulated insecticide designed for use in the control of exposed stages of stored product pests infesting warehouses containing non-perishable, packaged or bagged, raw or processed food commodities. The term "non-perishable" means any raw or processed food not subject to rapid decay or deterioration that would render it unfit for consumption.

Raw commodities would include cocoa beans, field-dried beans, grains, figs and nuts. Not included are: eggs, milk, meat, poultry, fresh fruits and vegetables.

Processed food includes: flour, sugar, cereals, packaged goods and crackers. Not included, are: hermetically sealed foods or manufactured dairy products and other processed foods requiring refrigeration.

TurboCide with DDVP, when applied as an aerosol, is recommended for uses in previously listed areas for the control of certain flying and crawling insects.

TurboCide with DDVP kills by contact, and the deposit of insecticide releases a vapor which continues to kill for a period of three or four days. TurboCide with DDVP vapor will give good control of flying insects, the exposed stages of crawling insects listed, beetles and young larvae, as well as destroying the viability of exposed eggs in the later stages of incubation.

Preparation of area to be treated: Clean up any debris prior to treatment. Shut off all air conditioners before using.

Unless directed otherwise (see Sanitary Sewers) apply TurboCide with DDVP at a rate of 5 grams per 1000 cubic feet of space (0.5 grams of DDVP) to control the listed pests in use areas shown on this label.

To reduce severe infestations of listed pests in warehouses use at a rate of 10 grams per 1000 cubic feet of space (1.0 gram of DDVP).

All occupants must be removed from dwelling or building before treatment is started and should not re-enter building or treated area for at least two hours. Open doors and windows and allow to be ventilated before occupying. Unpackaged or unwrapped food should be removed before treatment.

Use as an aerosol in edible products areas of food/feed processing areas only when plant is shut down. When applied as a contact spray, do not use in the edible product areas of food processing plants, restaurants, or other areas where food/feed is commercially prepared or served. Do not use in serving areas while food is exposed.

Food handling equipment, dishes and other equipment that contacts food should be covered with paper or washed with an effective cleaning compound followed by a potable water rinse after application. Do not contaminate water, feed or foodstuffs.

Remove all pets. Cover (or remove) fish tanks.

Cover or remove house plants within five feet of discharge to prevent damage to tender foliage.

006 44

006 03

BRAIN HARVESTING/HANDLING EQUIPMENT

TurboCide with DDVP is a convenient method of disinsecting harvesting machinery and all grain handling equipment prior to harvest. The principal requirement for this application is to ensure that all spaces in the machine receive an adequate dose of insecticide. Where access is readily available, TurboCide with DDVP should be sprayed into the space until the spray can be seen emerging from the furthest exit. Where there is no access to a space, a small hole should be drilled to admit the probe.

GREENHOUSES, LATHHOUSES, NURSERIES (HORTICULTURAL) AND SHADE HOUSES

Close all doors, windows and ventilators. Placard all entrances to prevent unprotected persons from entering the treated area. Be sure all foliage and blossoms are dry. Adjust inside temperature to 80 F \pm 5. Direct spray over, not at foliage.

MUSHROOM PROCESSING • MUSHROOM PRODUCTION (HOUSES)

For the control of adult Sciarid Flies and Phorid Flies, apply once every four days as a preventative or as insects appear. Repeat as necessary to maintain control. May be applied up to one day before harvest.

PEANUT WAREHOUSES

For the control of Almond Moth, and Indian-meal Moth, which are most active during the evening, application about sunset \pm 2 hours, will result in the most effective control.

SANITARY SEWERS (MANHOLES)

Cockroach and Mosquito infestations may be treated by inserting the extended nozzle through the vent hole in manhole covers and discharging 50 grams (5 grams of DDVP) of product. When treating sewers, one manhole per 200 feet of line should be treated.

TOBACCO FACTORIES, TOBACCO WAREHOUSES

To control the Tobacco Moth and Cigarette Beetle, which are most active during the evening, application about sunset (\pm 2 hours) will result in the most effective control.

NOTE TO PHYSICIAN

SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine should be injected at ten minute intervals in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. 2FAM is also antidotal and may be administered in conjunction with atropine. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by blood test.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: This cylinder must be checked periodically. A defective cylinder should not be used. A cylinder that has parts missing, is plainly worn, or is distorted, should be replaced immediately. Store upright in a cool, dry area.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide is a violation of Federal law. Pesticide that cannot be used according to label instructions must be returned to the point of purchase in its original container.

CONTAINER DISPOSITION: Cylinders must be returned to point of purchase for credit. All cylinders, valves and caps remain the property of the manufacturer. Federal regulations prohibit the refilling of cylinders without written consent of the manufacturer.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

USE AREAS

FOOD AND FEED PROCESSING AND HANDLING SITES

Bakeries, Beverage Plants, Bottling Plants, Breweries, Brewery Warehouses, Canning Plants, Egg Plants, Flour Mills, Food Handling Establishments, Food Processing Plants, Grain Elevators, Grain Mills, Granaries, Grocery Stores, Meat Packing Plants, Milk Rooms, Mushroom Production (Houses), Peanut Warehouses, Poultry Plants, Rabbit Processing Plants, Restaurants (Nonfood Area and Serving Areas), Rice Mills, Supermarkets and Wineries.

COMMERCIAL TRANSPORTATION FACILITIES

Aircraft, Boats, Buses, Freight Containers, Passenger Railroad Cars, Planes, Railroad Cars (Empty), Ships, Ships' Holds, Trains, Transportation Equipment, Trucks (Empty) and Truck Trailers.

PRIVATE OR COMMERCIAL DWELLINGS

Apartments, Cabins, Condominiums, Dwellings, Homes (Households), Hotels, Jails, Mobile Homes, Motels, Prisons, Recreational Vehicles and Shelters for Homeless.

ANIMAL SHELTERS

Animal Areas/Quarters, Cattle Barns, Dairy and Livestock Barns, Dog Kennels, Goat Houses, Horse Stables/Barns, Kennels, Loafing Sheds, Poultry Houses, Research Animal Quarters, Swine Houses and Zoos.

NONRESIDENTIAL BUILDINGS

Churches, City Missions, Drug Stores, Greenhouses, Lathhouses, Libraries, Mausoleums, Museums, Office Buildings, Shade Houses, Shopping Malls, Textile Mills and Warehouses, Theatres, Tobacco Factories and Warehouses, Utility Rooms, Voids, i.e.: Attics, False Ceilings, Walls and Equipment Voids of the above.

SCHOOLS AND DAY CARE CENTERS

Precautionary Note: Application is limited to only when these facilities are unoccupied. Keep children off treated surfaces until spray has dried.

HOSPICES, HOSPITALS AND NURSING HOMES

Precautionary Note: This product is a cholinesterase inhibitor and application is not allowed in patient rooms, therefore, application is limited to the unoccupied, nonpatient areas of these facilities.

MISCELLANEOUS SITES

Dumpsters, Garbage Compactors, Garbage Dumps, Grain Harvesting/Handling Equipment, Junk Yards, Manure Piles, Nurseries (Horticultural), Landscapes and Sanitary Sewers (Manholes).

SPACE AND CONTACT APPLICATIONS

To Control accessible adult states of:

Almond moths	Grain Mites
Angoumois grain moths	Granary Weevils
Ants	Hide Beetles
Bedbugs	Horn Flies
Brown Dog Ticks	Hornets
Cadellies	Horse Flies
Carpet beetles	House Flies
Centipedes	Indian-meal moths
Cereal beetles	Khapra Beetles
Cigarette beetles and young	Lesser Grain Borer
larvae; also destroying the	Lice
viability of eggs in the	Mediterranean Flour Moths
latter stages of incubation	Mosquitoes
if they are exposed	Mushroom Flies
Clothes moths and exposed eggs	Phorid Flies
Cockroaches	Sciariid Flies
Cocoa Bean Moths	Palmetto Bugs
Confused flour beetles	Red Flour Beetles
Crickets	Rice Weevils
Dark meal worms	Scorpions
Deer Flies	Silverfish
Dermestids	Small Flying Moths
Drugstore Beetles	other small flying
Earwigs	insects
Face Flies	Snails
Firebrats	Sowbugs
Fleas	Spiders
Flies	Stable Flies
Flour Beetles	Tabanids
Fly Maggots in maggot	Ticks (Brown Dog,
breeding areas	American dog,
Fruit Flies	Lone Star)
Fungus Gnats	Tobacco Moths
Gnats	Vinegar Flies
Grain Beetles	Wasps
Waterbugs	Mealy Bugs
Yellow Jackets	Omnivorous Leaf
Yellow Mealworms	Rollers
And The Following Green-	Red Spider Mites
house or Horticultural	Spider Mites
pests:	Thrips
Aphids	Vegetable Leaf
Cabbage Loopers	Miners
Cabbageworms	Whiteflies
Chrysanthemum Leaf	
Miners	