


12m 15

10F 3

4816-696
w345 / 160 295
17 4

Lawrence J. Feller
Fairfield American Corporation
8th and Harrison Streets
Frenchtown, NJ 08825

Dear Mr. Feller:

Subject: Amendment - Label Update
Permanone General Purpose Aqueous
Insecticide
EPA Registration No. 4816-696
Your Application Dated July 3, 1985

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

At the next label printing add the phrase "For Commercial/Industrial Use Only" on the front panel of the labeling.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

88064:LaRocca:RD-59:KENCO:10/11/85:10/24/85:ab:ps

CONCURRENCES

SYMBOL ▶	LaRocca						
SURNAME ▶							
DATE ▶							

BEST AVAILABLE COPY

PERMANONE® GENERAL PURPOSE AQUEOUS INSECTICIDE

2/85

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours. Ventilate before re-occupying.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from horn flies, face flies, house flies, mosquitoes and ticks. Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control stable flies, horse flies and deer flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control poultry lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control bedbugs and mites in poultry houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON PETS: To control fleas, ticks and lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

NOTE: Not recommended for use on kittens or puppies less than four weeks old.

OUTDOOR APPLICATION

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese beetles hit by spray as well as whiteflies, aphids, exposed thrips, spiders, red mites and leafhoppers.

May be used on plants, as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violers, asters, azaleas, begonias, carellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, euonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against whiteflies, aphids, spider and red mites, exposed thrips, lace bugs, Japanese beetles, pavement ants, clover mites and army worms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest (low, marshy places, stagnant pools, or ponds and shorelines). Apply at a rate of 2.5

pints per acre (0.9 ounce per 1000 square feet). Using truck-mounted conventional misting equipment such as Rotamist models SIC, 91 or 100 HT traveling at 5 m.p.h. to treat a 500 foot swath, calibrate equipment to deliver 0.95 gallon per minute. Under no conditions should open water be treated with this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Do not expose to freezing temperatures. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

CONTAINERS ONE GALLON AND SMALLER: Do not transport or store below 32°F. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contamination of feed and food-stuffs. Remove pets and birds, and cover fish aquaria before space spraying or surface applications.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one to two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, ponds, streams, tidal marshes or estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

BATCH

Printed in U.S.A.

BEST AVAILABLE COPY