

P.A 17 4816-697

Pg 183

MAR 31 1989

300	264405
17	1


Mr. Jan Brill
 Fairfield American Corporation
 809 Harrison Street
 Frenchtown NJ 08825

Dear Mr. Jan Brill

Subject: Rotacide EC
 EPA File Symbol/Reg. No. 4816-697
 Your submission dated February 28, 1989

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,



Phil Hutton
 Product Manager (17)
 Insecticide-Rodenticide Branch
 Registration Division (H7505C)

58831:I:A-4:Schroeder:C.Disk:KENCO:06/07/90:CL:VO:DD:CL

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

EVER - 2

ACCEPTED

ROTACIDE® EC

EPA Reg. No. 4816-697

EPA Est. No. 279-NY-1

- For use on vegetables and ornamentals (outdoors and in greenhouses, plantscapes and lathouses), forest, shade, fruit and nut trees.
- Designed especially for the "minor use" crop grower and "organic" farmers.
- Can be used up to and including day of harvest.

ACTIVE INGREDIENTS:

Rotenone	5.0%
*INERT INGREDIENTS	95.0%
	100.0%

*Contains Xylene Range Aromatic Solvent

PRECAUTIONAL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

TRANSLATION: TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.

ROTACIDE - Trademark of Fairfield American Corporation.

KEEP OUT OF REACH OF CHILDREN WARNING

Poisonous if swallowed or absorbed through skin
STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Contact a physician or Poison Control Center immediately.

FOR SKIN CONTACT: Wash immediately with soap and water.

IF IN EYES: Irrigate eyes with water. Contact a physician immediately.

NOTE TO PHYSICIAN: Gastric lavage, if ingested, followed by saline catharsis. No oils.

See Reverse Side for additional precautions



NET CONTENTS
 FAIRFIELD
 AMERICAN
 CORPORATION
 201 ROUTE 17 NORTH RUTHERFORD, NJ 07070
 72425 2/89

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried.

Apply at up to 0.4 lb active ingredient per acre and repeat as required to maintain effective control. Use in sufficient water for thorough coverage of upper and lower leaf surfaces.

To be tank mixed with BUTACIDE® 8 E.C. (or other 8 lb/gal. a.i. emulsifiable piperonyl butoxide) as recommended by your local Agricultural Extension Agent, consultant or pest control advisor, to control COLORADO POTATO BEETLES on POTATOES, EGGPLANTS AND TOMATOES. Tank mix at the rate of 1 part active Rotenone with 2 parts active Piperonyl Butoxide (76.8 fl. oz. of ROTACIDE® E.C. and 8.0 fl. oz. of BUTACIDE 8 E.C.). This will provide 0.25 lb. of Rotenone and 0.5 lb. of Piperonyl Butoxide per acre.

COMBINED WITH OTHER INSECTICIDES:

May be tank-mixed with other insecticides, as recommended by your local Agricultural Extension Agent, Crop Consultant or Pest Control Advisor, to control the insects on the crops named on their labels. Observe the directions on the insecticide label, tank-mix at a rate of up to 0.4 lb A/A of Rotacide® EC (to provide 0.4 lb of active ingredient, use 1 gallon of Rotacide® EC.) Tank-mix applications should always be made in accordance with the more restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

USE THROUGH IRRIGATION SYSTEMS (CHEMIGATION):

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the pump motor stops. The irrigation line

or water pump must include a stop the water pump motor when the pressure decreases to the point where the system is severely affected. Systems must be equipped with a positive displacement injection pump effectively designed and tested to be compatible with pesticides. The system must have a system interlock. Do not use systems that favor drift beyond the area in which the pesticide is being applied. Accuracy in calibration and distribution of the pump is essential. Projecting a larger volume of moisture may result from excessive overlap of sprinkler distribution patterns.

NOTE: MAKE APPLICATIONS AS EARLY AS POSSIBLE

Rotacide is relatively nontoxic to humans. This benefit apply early in the season.

GROWING CROPS (OUTDOOR HOUSES):

ROOT AND TUBER VEGETABLES: limited to, Arracacha; arrowroot; burdock, edible; carrot; cassava (cassia root); chervil, turniproot (taro); ginger; ginseng; horseradish; noprooted; parsnip; potato; radish (Daikon); rutabaga; salsify (Oxalis); Spanish; skirret; sweet potato; turmeric; turnip; yam, true; yam.

BULB VEGETABLES (Allium): limited to, Garlic; leek; onion;

LEAFY VEGETABLES: (Including Amaranth (leafy amaranth, Callalota); rugula (Roquette); celery; celeriac; chrysanthemum, edible leafy vegetable; cress, garden; cress, Upland (Dandelion); dock (sorrel); endive; lettuce; orach; parsley; purslane; rhubarb; spinach; spinach, fine-leafed; New Zealand; swiss chard).

BRASSICA (cole) LEAFY VEGETABLES: not limited to, Broccoli; broccolo; raab (rapini); Brussels sprout (bok choy); cabbage, Chinese; cauliflower; collards; kale; kohlrabi; greens).

CONTINUE