4816-7		06/30/	11999	Form Appro	OMB No. 207	1/10 0-0060. Approval expires 2-28-95	
≎ EPA	Environmental	nited States	Agency		Registration Amendme Other	OPP Identifier Number	
Application for Pesticide - Section 1							
1. Company/Product Number 4816-707 4. Company/Product (Name) Kicker			2. EPA Product Manager Marien Johnson PM#			3. Proposed Classification None Restricted	
5. Name and Address of Applicant Include ZIP Codel Agrevo Environmental Health 95 Chestant 12: dge 120a d Montvalt, NJ 07645 Check if this is a new address			6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name				
		S	Section -	11	<u> </u>	NOTIFICATION:	
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.				Final printed labels in response to Agency letter dated "Me Too" Application. Other - Explain below.			
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Notification of label change per PR Notice 96-6. This notification is consistent with the provisions of PR Notices 96-6 and 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notices 96-6 and 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. NOTIFICATION OF REVISED WARRANTY STATEMENT PER PR NOTICE 98-10 AND REVISED STATEMENTS REGARDING PET USES PER PR NOTICE 96-6.							
		S	ection -				
1. Material This Produc Child-Resistant Packagi Yes* No * Certification must be submitted	Unit Packaging Yes No	No per If	Yes No "Yes" ackage wgt	Packaging No. per container	P	ntainer Metal Plastic Glass Paper Other (Specify)	
Location of Net Cont Label	ents Information Container	4. Size(s) Retail Co	ontainer	5. Lo	On Label I	Directions accompanying product	
6. Manner in Which Label is Affixed to Product Lithograp Paper glu Stenciled			Other				
	Section - IV						
1. Contact Point (Com	plete items directly below	for identification of	individu al to	be contacted, if nec	essary, to proce	ess this application.)	
Name Sharon M. Johnston			Title Telephone Nc. (Include Area Code) Manager, Refistrations (201) 307-3363				
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and councilete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.							
2. Signature		3. Ti	tle				
	5	m	lanager.	Registrat	s		
4. Typed Name	r. Johnston	5. D	ate	: 15 1999			
U TREOR A	I, JOHNSTON		J W ~ 1	-,			



6.0% 60.0%

ACTIVE INGREDIENTS:
Pyrethrins
*Piperonyl Butoxide, Technical
**OTHER INGREDIENTS

34.0%

100.0%

*Equivalent to min. 48.0% (butylcarbityl)(6-propylpiperonyl) ether and 12.0% related compounds. **Contains Petroleum Distillate

Kicker, Pyrenone & Actisol - Registered Trademarks of AgrEvo Environmental Health, Inc.

KEEP OUT OF REACH OF CHILDREN CAUTION SEE BOOKLET FOR FIRST AID AND PRECAUTIONARY STATEMENTS

Net Contents: 1 pint

5/99

NOTIFICATION

READ ENTIRE LABEL BEFORE EACH USE

FOR PET USE: USE ONLY ON DOGS AND CATS

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

See Booklet for Complete Directions For Use

STORAGE & DISPOSAL
Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool dry place inaccessible to children and pets.

Disposal: Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

AgrEvo Environmental Health

95 Chestnut Ridge Road Montvale NJ 07645

EPA Reg. No. 4816-707

15303

EPA Est. No.: 10370-TX-1

5/99

FIRST AID

FIRST AID

If SWALLOWED: Contact a physician or Poison
Control Center immediately. Do not induce
vomiting unless directed by a physician. This
product contains periodicum disulfate and vomiting
may cause aspiration pneumonia.

If IN EYES: Flush with water for approximately 15
minutes. Contact a physician d irritation persists.

If ON SKIN: Remove contaminated cipthing, Wash
skin with plants of sour and water. Consert. skin with plenty of soap and water. Contact a physician if irritation persists.

If INHALED: Remove victim to fresh air, Apply artificial respiration it indicated.

In case of Medical emergencies or health and safety inquiries or in case of fire, leaking or damaged containers, information may be obtained by calling 1-800-471-0660.

For product information Call Toll-Free: 1-800-331-2867

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

CAUTION

Harmful if swallowed. Do not get in eyes, Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not spray directly infon eyes, mouth or genitalia of pets. Do not use on dogs or cats under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using any pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rines with large amounts of water. If signs continue, consult a veterinarian immediately.

When using this product as a space spray in food processing plants, foods should be removed or

covered during treatment, Do not apply as a spray while food processing is underway, Food processing surfaces and equipment must be covered during treatment or cleaned with a suitable detergent and rinsed with potable water before reuse. Except in Federally inspected meat and poultry plants, lood processing operations may continue when this product is applied as a surface spray with care and in accordance with the directions and precautions given

Environmental Hazards

Environmental Hazards
This product is toxic to tish, shrimp, crab and other
aquatic organisms. Do not apply directly to water, to
areas where surface water is present, or to interridal
areas below the mean high watermark. Shrimp and
crab may he killed all application rates recommended
on this label. Do not apply where runoff is likely to
occur. Do not apply may be a produced to the conditions tavor
drift from areas treated.

Physical and Chemical Hazards **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsisient with its labeling. Not for use on turl being grown for sale or other commercial use as sod, or for commercial seed commercial use as sod, or for commercial seed production, or for research purposes. A highly versalile Pyrenone Insecticide Concentrate for indoor or outdoor application as a surface, space, area or confact spray. Kicker may be applied through mechanical, serosol or ULV generators as well as thermal fogging equipment and conventional fogging or spraying equipment. Kicker may be applied as a general surface, space, spot or crack and crevice treatment in both food and non-food areas of food handling establishments and is authorized for use in all areas of official establishments operating under the Meat. Poultry, Shell Egg Grading and Egg Products Inspection Programs.

Kicker is intended for use in and around buildings and structures as well as modes of transport. INDOOR USE AREAS: Apartment Buildings, Bakeries, Beverage Plants, Bottling Facilities.

Breweries, Cafeterias, Candy Plants, Canneries, Cereal Processing and Manufacturing Plants, Dairies and Dairy Product Processing Plants, Frozen Food Plants, Grin Mills, Granaries, Homes, Hospitals, Hotels, Houses, Industrial Buildings, Kennels, Kitchens, Laboratories, Manufacturing Facilities, Material Buildings, Restaurants, Mosels, Spice Plants, Med and Vegetable Canneries, Motels, Nursing Homes, Office Buildings, Restaurants, Schools, Spice Plants, Stores, Supermarkets, Warehouses, Wineries and Similar Structures. OUTDOOR AREAS: Drive-in Restaurants, Drive-in Theaters, Golf Courses, Parks, Plangrounds, Recreational Areas, Urban Areas and Zoos.
MODES OF TRANSPORT: Butes, Trucks, Trailers, Rail Cars and Marine Vessels.

٠.;

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have direct. The insecticidal components of Kicker, pyrethrism and piperonyl butoxide, are classified as food additives and established tolerances will not be exceeded when used as directed. Kicker may be difuted with water or oil and applied with conventional application equipment including, but not limited to. compressed air sprayers, power sprayers, mechanical aerosol generators (ULV generators or cold foggers) and thermal foggers foil dilutions only). When diluting with water, prepare only the amount needed for immediate use and agitate occasionally during use. Holding a water dilution more than 24 hours is not recommended, for oil dilutions that are applied in food handling establishments, a deodorized petroleum oil which conforms with CFR 21 (especially section 172.884) is required (such as Exxon isopar M, Ashland Low Odor Base Solvent or their equivalents).

Tank-mix applications must be made in accordance with the more restrictive oil label limitations and precautions. No label dosages may be exceeded. This cannot be tank-mixed with any product with

label prohibitions against such mixing.
INDOOR USE AS A CONTACT AND SURFACE SPRAY

GENERAL PEST CONTROL: To control accessible, exposed stages or crawling insects including, but not limited to. Ants. Coctroaches, Cadelles, Cigarette Beetlea, Confused Flour Beetles, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Grain Mites, Red Flour Beetles, Rice Weevils, Saviotothed Grain Beetles, Spider Beetles, Fellow Mealworms in sites that include, but are not limited to, apartments, homes, restaurants, food processing plants, industrial installations and warehouses, dilute at the rate of up to 4.25 fluid ounces of Kicker per gallom of water and apply at the rate of 1 gallon to 750 square feet. Thoroughly spray the surfaces where these insects are usually found including walls, basehoards, moldings, doors, cabinets, shelves, sinks and pipes with special attention to cracks and revices, niches, dark corners, drains and other harborage sites.

Except in Federally inspected meat and poulity plants, for the property of the processing operations may continue when Kicker is applied as a surface, spot or crack and crevice treatment in accordance with the directions and precaulisms given above.

DILUTION CHART

Fluid Ounces	Approximate				
Of Kicker Per Gallon	% Pyrethrins	% Piperonyl Butoxide			
0.25	0.0125	0.125			
0.5	0.025	0.25			
1.0	0.05	0.5			
2.125	0.1	1.0			
4.25	0.2	2.0			

IN USDA INSPECTED FACILITIES: To control accessible, exposed stages of crawling insects including, but not limited to, Ants, Cockroaches, Cadelles, Cigarette Beetles, Confused Flour Beetles, Dark Mealwoms, Dried Fruit Beetles, Drugstore Beetles, Grain Mites, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles, Spider

Beetles, Yellow Mealworms, dilute 2.125 Iluid ounces of Kicker per gallon of water and apply at the rate of 1 gallon to 750 square teet. Thoroughly spray the surfaces where these insects are usually found including walls, baseboards, moltings, doors, cabinets, shelves, sinks and pipes with special attention to cracks and crevices, niches, dark comers, drains and other harborage sites.

IN COMBINATION WITH RESIDUAL

INSECTICIDES (including, but not limited to, Baygon, Commodore, Conquer, Cynoff, Demon, Diazinon, Dursban, Empire, Ficam, Pyrid, Satrotin, and Tempol:

To provide flushing and quick knockdown of insects: Kickor may be tank-mixed with other insecticides at the rate of % to % ounce (equivalent to % to 1 tablespoon or 7.4 to 14.8ml) per gallon of finished soray.

To provide increased control of resistant German cockroaches: tank-mix 2.125 to 4.25 fluid ounces of kicker with the appropriate amount of companion insecticide per finished gallon and apply per the Directions for Use on the companion insecticide label, in USDA inspected facilities, tank-mix 2.125 fluid ounces of Kicker with the appropriate amount of companion insecticide per finished gallon and apply per the Directions for Use on the companion insecticide label.

Prior to tank-mixing, a small jar compatibility test should be conducted using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture. When tank-mixing Kicker with wettable powder to mulations, tirst add the wettable powders to the diluent and agitate to thoroughly suspend it in solution and then add the desired amount of Kicker to the diluted wettable powder. Tank-mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions sagins such mixing.

against south imanig.
Baygon, Commodore, Conquer, Cynoff, Demon,
Dursban, Empire, Ficam, Pyrid, Salrotin and Tempo

are Registered Trademarks of Bayer, Zeneca Inc., Paragon Professional Pest Control Products, FMC Corporation, DowElanco, AgrEvo UK Limited, Terminix International Inc., and Sandoz Ltd., respectively.

FLÉA AND TICK CONTROL IN BUILDINGS: Dilute 2.125 to 4.25 fluid ounces of Kicker per gallon of water and apply the spray throughly to infested areas such as pet beds and resting quarters, nearby cracks and crevices, rugs and floor coverings (including under their edges) between and under the cushions of upholstered furmiture and other areas where these pests may be present. Entire rugs and turniture should be treated at the rate of 1 gallon of dilute spray per 750 square feet of surface. In combination with conventional residual insecticides: To provide quick knockdown of adult fleas, trank-mix. O, 5 to 1.0 fluid ounces of Kicker with the appropriate amount of companion insecticide per finished gallon and apply per the Directions for Use on the companion insecticide

Prior to tank-mixing, a small jar compatibility test should be conducted using the proper proportions of chemicals and water to ensure the physical compatibility or the mixture. Tank-mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing, in combination with insect Growth Regulators: To provide quick knockdown of adult (leas, tank-mix 2.125 to 4.25 fluid ounces of Kicker with the appropriate amount of companion insecticide per finished gallon and apply per the Directions for Use on the companion insecticide label. Prior to tank-mixing, a small jar compatibility test should be conducted using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture. Tank-mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This

product carrick be mixed with any product with label prohibitions against such mixing. Per hedding should be removed and replaced with tresh bedding after treatment of pet areas. The pets should also be treated with an EPA approved product before they are allowed to enter the treated area. Dogs and cats may be treated with Kicker by diffusing 2.125 fluid ounces per gallon of water and spraying or sponging thoroughly to penetrate the hair and contact the skint to all parts of the animals body except the eyes, nose, mouth and genitalia.

PANTRY PEST CONTROL: To control exposed

nair and contact the skini to all parts of the animals hody except the eyes, nose, mouth and genitalia. PANTRY PEST CONTROL: To control exposed adult and immature stages of Stored Product insects including, but not limited to. Almond Moths, Angournois Grain Moths, Cadelles, Cigarette Beetles, Confused Flour Beetles, Orangstore Beetles, Flat Grain Beetles, Grains Weevils, Indianmeal Moths, Lesser Grain Borers, Maize Weevils, Mediterranear Flour Medite, Merchant Grain Beetles, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Sawtonhed Grain Beetles and Squarenecked Grain Beetles, dilute 2:125 to 4:25 tluid ounces of Kicker per gallon of water and apply to cupboards, shelves and storage areas. Remove all utensils, unprotected Goodstuffs for any with original packaging that has been opened), shelf paper and other objects before ireatment. Allow treated surfaces to dry and then cover shelves with clean paper before replacing any utensils, loodstuffs or other items. Discard any toxistuff that was accidentally contaminated with the spray solution.

PREMISE PEST CONTROL: Termite swarmers: To

the spray solution.

PREMISE PEST CONTROL: Termite swarmers: To control swarming termiles, dilute 2.125 to 4.25 fluid dunces of Kicker per gallon of water and apply as a surface stray at the rate of 1 gallon per 750 square leet. Apply as a crack and crevice treatment to the locations from which the termites are emerging.

to the locations from which the fermions are emerging.

Carpet Beetles: To control carpet beetles (black, furniture and varied), dilute 4.25 fluid ounces of Kicker per gallon of water and apply to the entire carpet at the rate of 1 gallon per 750 square feet.

Make localized applications under the edge of

carpets, floor coverings and furniture as well as along baseboards, in closets and to infested areas of shelving.

shelving. Bed Bugs: To control **Bedbugs**, citiute 2,125 fluid ounces of Kicker per gallon of water and apply thoroughly to the top, ends, sides and bottoms of mattresses paying special attention to lutis, seams and tolds. Treat bedsprings and frames, including all cracks and joints, Apply as a crack and crevice treatment to all baseboards, moldings, flooring, closets, shelves, furniture and picture frames that may provide harborage to bedsbugs. Webbing Clothes Moths: To control **Clothes Moths** remove miested articles from storage, brush them thoroughly and air-dry them in sunlight for several hours. Dilute 2.125 fluid ounces of Kicker per gallon of water and apply as a crack and crevice

gallon of water and apply as a crack and crevice treatment to empty chests, bureaus and storage sites. After airing, infested articles may be lightly treated with the spray solution (keep the nozzle at least three feet from the fabric to avoid staining).

FOR INDOOR APPLICATIONS: Remove pets (including birds) from the area being treated and cover fish aquaria before spraying.

cover fish aquaria before spraying.

INDOOR USE AS A SPACE SPRAY
TO control crawling and flying insects in sites that include, but are not limited to, food processing plants, IINCLUDING USDA INSPECTED
FACILITIES, inclustrial installations and warehouses, Kicker may be diluted with water and applied as a space spray. For best results, close doors and windows before spraying and keep them closed for 30 minutes after treatment. The applicator must vacate the area after treatment and ventilate before reoccupying, Where oil residues are not undesirable. Kicker can be diluted in decodorized base oil instead of water and applicators.

The use of this product in food processing or food with mechanical, inermal or ULY applicators. The use of this product in food processing or food handling establishments must be confined to time periods when the plant is not in operation. Foods must be covered or removed during treatment. Food processing surfaces and equipment must be covered during treatment or cleaned with a suitable

detergent and rinsed with potable water before

CRAWLING AND FLYING INSECTS: For control of accessible, exposed stages of CRAWLING INSECTS including, but not limited to, Ants, Cockroaches, Cadelles, Cigarette Beetles, Controsed Flour Beetles, Dark Meahworms, Dried Fruit Beetles, Drugstore Beetles, Grain Mites, Red Flour Beetles, Rice Weevils, Saw toothed Grain Beetles, Spider Beetles, Wellow Meahworms and FLYING INSECTS including, but not limited to, Angournois Grain Moths, Cheece Shippers, Fruit Flies, Fungus Gnals, Gnats, House Files, Indianneal Moths, Mosquitoes, Mediterranean Flour Moths, Small Flying Moths, Tobacco Moths, dilute 10,67 fluid ounces of Kicker per gallon of water or oil and apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spiral towards the ceiling and upper corners of the area and keep the area crisced for at least 30 minutes after treatment. Ventilate the CRAWLING AND FLYING INSECTS: For control of at least 30 minutes after treatment. Ventilate the area before reoccupying. Repeat treatment as necessary.

FLYING INSECTS: For control of accessible FAYING INSECTS: For control of accessible, exposed stages of flying insects including, but not limited to. Angaumois Grain Morits, Cheese Skippers, Fruit Files, Fungus Grats, Grats, House Files, Indiametal Moth, Mosquitoes, Mediterranean Flour Mether, Small Flying Moths, Tobacco Moths, dilute 2.67 fluid ounces of Kicker per gallon of water or oil and apply at the rate of 1 ounceper 1000 cubic feet of space. Direct the stray towards the ceiting and upper comes of the area and behind obstructions. Vacate the treated area and keep area closed for at least 30 minutes after treatment. Ventilate the area before reoccupying. Repeat treatment as necessary. VOID AREA TREATMENT: Undiluted ULV not limited to, attics, talse ceilings, wall and equipment voids, calculate the volume of the void and apply Kicker through Actisoft or other ULV crack and crevice application equipment at the rat of 0.33 fluid ounces per 1000 cubic feet of space.

color: black text

OUTDOOR USE
PERIMETER TREATMENTS: To control Ants, Bees, PERIMETER TREATMENTS: To control Ants, Bees, Centipedes, Cockroaches, Crickets, Firebrats, Fleas, Ground Beetles, Millipedes, Pillbugs, Silverfish, Sowbugs, Spiders, Ticks and Waps, dilute 2,125 to 4,25 thid ounces of Kicker per gallon of water and apply at the rate of 1 gallon per 750 square feet to a band of soil and vegetation to to 10 ieet wide around and adjacent to buildings. Treat the building foundation to a height of 2 to 3 feet at the rate of 1 gallon per 750 square feet.

the rate of 1 gallon per 750 square feet, TICK CONTROL: To control Brown Dog Ticks around residential areas, rural fields, woods, lots, roadsides, pathways and coastal vegetation areas, cliute 4,25 thuid ounces of Kicker per gallon of water and apply with a miss blower for other similar equipment calibrated to relieve 1 gallons per acre. Apply during the peak of infestation and repeat as necessary to maintain control. Thorough applications should be made to the vegetation upon which the ticks rest.

FLEA CONTROL: To comrol fleas around residential FILEA CONTROL: 10 Control Heas around residential areas, dop houses, kennels, paths, patios and under plants, shrubs, trees or other shaded, moist areas where pets may rest, dilute 4.25 fluid ounces of Kicker per gallon of water and apply at the rate of 1 gallon per 750 square feet.

In combination with other insecticides: To provide quick knockdown of the outdoor pests listed above when used with a residual insecticide, tank-mix 1.0 to 2.125 fluid ounces of Kicker with the proper amount of companion insecticite and apply per the Directions for Use on the companion insecticide label.

label. Prior to tank-mixing, a small jar compatibility test should be conducted using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture. Tank-mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

USE ON TURF AND GRASS: Used alone as a USE ON TURE AND GRASS: Used alone as a contacts pay 10 control Ants, Armyworms, Billbugs, Chinch Bugs, Chiggars, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, Hyperodes Weevils (adults), Japanese Beetles (adults), Mole Crickets, Sod Webworms And Ticks, dilute and apply per the instructions in the following table:

Treatment Ansa (Square Feet)	Fluid Ounces Of Kicker	Suggested Volume Of Water (Gallons)		
1000	0.25 to 0.5	2.5 to 5.0		
5000	1.25 to 2.5	12.5 to 25.0		
20000	5.0 to 10.0	50.0 to 100.0		
43560*	12.0 to 24.0	110.0 to 220.0		

a Dilute with enough water to obtain thorough coverage. b 43560 square feet = 1 acre.

to no you square test is later. In combination with other insecticides: To provide quick knockdown of insects when used with a residual insecticide, tank-mix Kicker at the tale listed above with the proper amount of companion insecticide and apply per the Directions for Use on the companion insecticide label.

the companion insecticide tabel.

Prior to tank-mixing, a small jar compatibility test should be conducted using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture. Tank-mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. This product cannot be mixed with any product with label prohibitions against such mixing.

As a rurf pest diagnostic aid: To detect turf insects

label prohibitions against such mixing.
As a rurf pest diagnostic aid: To detect rurf insects prior to making an insecticacle application or to evaluate control from previous treatments, dilute 0.5 fluid ounces of Kicker per gallon of water and apply evenly with a sprinking can over one square yard of turf. Record the species and number of insects present ten minutes after application. Sample 3 to 5 sites per S000 square teet. Note: this procedure does not bring white grubs or billbug grubs to the surface. Use other methods to sample

tor these pests.

tor these years.

USE AS A PET SPRAY
TO CONTROL FLEAS AND TICKS ON PETS: Dogs
and cats may be treated with Kicker by diluting at the
rate of 1.0 to 2.125 fluod ounces per gallon of water
and suraying or sponging thoroughly the penetrate the
hair and contact the skini to all pans of the animals
havly except the eyes, note, mouth and genitalia.

REPEAT APPLICATION EVERY TWO WEEKS, IF,
NECESSARY for best results against tleas and ticks on
flogs and cats, the animal quarters should also be
troated and bedding should be removed and replaced
after the per areas are treated.

after the pel areas are treated.

STORAGE AND DISPOSAL

FOR CONTAINERS LARGER THAN ONE GALLON: Do not contaminate water, food or

GALLON: Do not contaminate water, food or feed by storage or disjoinal.
PESTICIDE STORAGE AND SPILL.
PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdur, earth, fuller's earth, etc. Dispose of with chemical waste.

with chemical wase.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by

label instructions must be disposed or all or by an approved washe disposal facility. CONTAINER DISPOSAL: Triple rinse (or equivalent) them ofter for recycling or reconditioning, or puncture and dispose of in a saniary landfill, or by other approved State and local approximant.

saniary landing or by orner approved state local procedures. FOR CONTAINERS ONE GALLON AND SMALLER: STORAGE: Store in original container in a cool, dry place maccessible to children and pets.

DISPOSAL: Do not reuse container. Wrap

container in several layers of newspaper and discard in trash.

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the

following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed

product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond Agrevo Environmental Health's control, it is impossible for Agrevo Environmental Health to eliminate all risks associated with the use of this product. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED. OF MERCHANTABLITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Agrevo Environmental Health is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Agrevo

those contained herein or to modify the warranties contained herein. Agric to Environmental Health disclaims any fiability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), fort, negligence, strict liability or otherwise.

OINMANIER.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TOR, NECLICIENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT AGREVO ENVIRONMENTAL HEALTH'S ELECTION. THE REPLACEMENT OF PRODUCT.

OAgrévo Environmental Health, Inc., 1999