

BEST AVAILABLE COPY**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where food may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before reuse. Exposed food should be covered or removed.

Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches.

Remove birds and cover fish aquariums before spraying.

Houseflies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Blow Flies and Small Flying Moths: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Repeat treatment as necessary.

Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs and Pillbugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moths, Angoumois Grain Moths: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application as necessary.

Exposed stages of Confused Flour Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45

minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary. Allow all sprayed articles to dry thoroughly before use.

To control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining. Allow all sprayed articles to dry thoroughly before use.

In animal quarters and milk rooms: For rapid control of *House Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees and Small Flying Moths*, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

To treat livestock: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of *Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes, and Gnats:* Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

For Swine: Do not ship animals for slaughter within 5 days of last treatment.

To control Blood Sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control Poultry Lice, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of Bedbugs and Mites in Poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

SEE REVERSE

PERMANONE**MULTI-PURPOSE 0.25% SPRAY**

EPA REG. NO. 4816-714

EPA EST. NO. 279-NY-1

- * CONTAINS THE SYNTHETIC PYRETHROID... PERMANONE
- * CONTROLS RESIDENTIAL, INDUSTRIAL AND GARDEN INSECTS
- * FOR USE ON DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS
- * A Water-Based Aerosol
- * CONTROLS NAMED PESTS FOR UP TO FOUR WEEKS

ACTIVE INGREDIENT:

*Permethrin.....0.25%

INERT INGREDIENTS:.....99.75%

100.00%

*(3-phenoxyphenyl)methyl (+/-) cis/trans 3-(2,2-dichloroethyl) 2,2-dimethylcyclopropanecarboxylate

Cis/Trans ratio: Min. 35% (+/-) cis and max. 65% (+/-) trans.

PERMANONE - Registered Trademark of Fairfield American Corporation

KEEP OUT OF REACH OF CHILDREN

CAUTION**STATEMENT OF PRACTICAL TREATMENT**

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

IF IN EYES: Flush with plenty of water. Contact a physician if irritation persists.

IF ON SKIN: Wash affected areas immediately with soap and water. Get medical attention if irritation persists.

See reverse for additional precautionary statements

NET CONTENTS



FAIRFIELD
AMERICAN
CORPORATION

201 ROUTE 17 NORTH, RUTHERFORD, NJ 07070

80020 112189F



PROPERTY OF THE U.S. GOVERNMENT
NOT TO BE REPRODUCED OR
TRANSMITTED IN ANY FORM
OR BY ANY MEANS
WITHOUT PERMISSION

- **Fleas and Ticks: (Adults and Larvae):** Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floorcovering with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

To control **Fleas, Ticks and Lice on Dogs.** Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat application weekly if necessary to control infestations. NOTE: Not recommended for use on puppies less than four weeks old.

To kill **Lice and Louse eggs:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches.

Spray each square foot for three seconds.

Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned.

Allow all sprayed articles to dry thoroughly before reuse.

For the Control of Garden Insects: For use on roses, dahlias, asters and other ornamentals to kill *Japanese Beetles* hit by spray, as well as *Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites and Leafminers.* May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas. May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias. Effective when used as directed against *Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms.* When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

STORAGE AND DISPOSAL

Store in a cool, dry area. Do not transport or store below 32°F. Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Avoid spraying in eyes. Avoid breathing spray mist. Avoid contact with skin. Wash hands thoroughly after use. Vacate room after treatment and ventilate before reoccupying. Avoid contamination of feed and foodstuffs. Cover or remove fishbowls. This product is not for use on humans.

Animals: Do not spray directly in/on eyes, mouth or genitalia of pets. Do not treat or cause exposure to kittens or puppies less than four weeks old.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

BEST AVAILABLE COPY