

Reg # 4816-714

PM-13

124

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460  <b>NOTICE OF PESTICIDE:</b> <input type="checkbox"/> REGISTRATION <input checked="" type="checkbox"/> REREГИSTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	EPA REGISTRATION 4816-714	DATE OF ISSUANCE NOV 15 1992
	TERM OF ISSUANCE Until Reregistered	
	NAME OF PESTICIDE PRODUCT Permanone Multi-Purpose 0.25% Spray	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Fairfield American Corporation  
809 Harrison Street  
Frenchtown, NJ 08825

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

- a. Add the phrase "EPA Registration No. 4816-714."
- b. Add NAMED after CONTROLS in "Controls pests for up to four weeks."
- c. Change ACTIVE INGREDIENTS to the singular.
- d. Change "See rear panel for additional precautions" to "See rear panel for additional precautionary statements."

**BEST AVAILABLE COPY**

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

*[Handwritten Signature]*

DATE

11/15/92

- e. Since your data matrix does not give a dielectrical breakdown voltage, you must add the following to the use directions:

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

- f. Expand the statement beginning "Do not use in the edible . . ." to read:

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where food may be prepared or held. In the home all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before reuse. Exposed food should be covered or removed.

Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

- g. Use in food-handling establishments can imply use in U.S. Department of Agriculture (USDA) Meat and Poultry Plants. Therefore, either add the statement "Not for Use in USDA Meat and Poultry Plants," or submit an application to USDA requesting use in these areas. If granted, you can then indicate on the label that the product can be used in these areas.
- h. Under "Bed bugs" and "Clothes Moths" directions, add the following precaution "Allow all sprayed articles to dry thoroughly before use."

3. The 30-day elevated storage stability data are acceptable; however, please provide a 1-year room temperature study.

4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

**BEST AVAILABLE COPY**

CODE

# PERMANONE® MULTI-PURPOSE 0.25% SPRAY

EPA REG. NO. 4816-

EPA EST. NO. 279-NY-1

CONTAINS THE SYNTHETIC PYRETHROID ...

PERMANONE®

CONTROLS RESIDENTIAL, INDUSTRIAL AND GARDEN INSECTS

FOR USE ON DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS

AN AQUEOUS, STABLE, MICRO EMULSION

CONTROLS PESTS FOR UP TO FOUR WEEKS

**ACTIVE INGREDIENTS:**

\*Permethrin ..... 0.25%

**INERT INGREDIENTS:** ..... 99.75%

100.00%

\*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2, 2-dimethylcyclopropanecarboxylate Cis/Trans ratio: Min. 35% (+/-) cis and max. 65% (+/-) trans.

PERMANONE - Registered Trademark of Fairfield American Corporation

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

See rear panel for additional precautions.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use in the edible product areas of food/feed processing plants, restaurants or other areas where food/feed is commercially prepared or served. Do not use in serving areas while food is exposed.

**SHAKE WELL BEFORE USING**

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches.

Remove birds and cover fish aquariums before spraying.

**Houseflies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Blow Flies and Small Flying Moths:** Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Repeat treatment as necessary.

**Cockroaches, Waterbugs, Palmello Bugs, Silverfish, Firebrats, C. Scorpions, Spiders, Millipedes, Centipedes, Sowbugs, and P.** Spray into hiding places such as cracks and crevices, behind cabinets, along baseboards and floors, around drains and plumbing. Hitting insects with spray whenever possible. Repeat treatment as necessary.

**Ants:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

**Indian Meal Moths, Angoumols Grain Moths:** Close doors and windows. Direct spray into all parts of room especially around stored food products, containers, pallets and darkened areas. Spray 7 to 10 seconds for 1000 cubic feet of space. Keep area closed for 15 minutes for treatment. Repeat application as necessary.

**Exposed stages of Confused Flour Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites:** Spray into cracks and crevices, processing and handling equipment, around containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

**Carpet Beetles:** Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

**Bedbugs:** Spray mattresses, particularly around tufts and seams. Separate beds apart and spray into all joints. Treat baseboards, molding and floors. Repeat treatment as necessary.

**To control Clothes Moths:** Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Spray liberally to empty chests, closets, bureaus and other storage areas, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

**In animal quarters and milk rooms:** For rapid control of Houseflies, Mosquitoes, Gnats, Wasps, Hornets, Bees and Small Flying Moths, spray windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is complete.

NET CONTENTS: \_\_\_\_\_

**BEST AVAILABLE COPY**

SEE REVERSE SIDE

484

Effective when used as directed against Whiteflies, Aphids, Spider and Red exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat as infestations occur.

### STORAGE AND DISPOSAL

Store in a cool, dry area. Do not transport or store below 32° F.

Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

Avoid spraying in eyes. Avoid breathing spray mist. Avoid contact with skin. Wash hands thoroughly after use. Vacate room after treatment and ventilate before reoccupying. Avoid contamination of feed and foodstuffs. Cover or remove fish bowls. This product is not for use on humans. If lice infestations should occur on humans, consult either your physician or pharmacist for a product for use on humans.

Animals: Do not spray directly in/on eyes, mouth or genitalia. Do not cause exposure to puppies less than four weeks old.

#### STATEMENT OF PRACTICAL TREATMENT

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

IF IN EYES: Flush with plenty of water. Contact a physician if irritation persists.

IF ON SKIN: Wash affected areas immediately with soap and water. Get medical attention if irritation persists.

#### PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**BEST AVAILABLE COPY**

To control Blood Sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control Poultry Lice, spray roost walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of Bedbugs and Mites in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

Fleas and Ticks: (Adults and Larvae): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

To control Fleas, Ticks and Lice on Dogs. Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat application weekly if necessary to control infestations. NOTE: Not recommended for use on puppies less than four weeks old.

To kill Lice and Louse eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 3 to 10 inches.

Spray each square foot for three seconds.

Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned.

Allow all sprayed articles to dry thoroughly before use.

For the Control of Garden Insects: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites and Leafminers.

May be used on plants, as well as a multi-purpose spray in small gardens for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.