

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460  <b>NOTICE OF PESTICIDE:</b> <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended)	EPA REGISTRATION NO.	DATE OF ISSUANCE
	4816-745	JUL 14 1982
	TERM OF ISSUANCE Until Reregistration	
NAME OF PESTICIDE PRODUCT PyraPerm Insect Kill WBA N67A		

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Fairfield American Corporation  
 809 Harrison Street  
 Frenchtown, NJ 08825

191 / 4 / 2272

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c) (7) (A) since you have agreed to:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA sec. 3(c) (5) or FIFRA sec. 4 when the Agency requires all registrants of similar products to submit such data.

2. Submit data listed below, conducted in accordance with the Pesticide Assessment Guideline (40 CFR Part 158 - Data Requirements for Registration) no later than one year from receipt of the Notice for Registration.

a. Dermal Sensitization - Guideline 81-6

3. Make the labeling changes listed below before you release the product for shipment:

- a. Add the phrase, "EPA Registration No. 4816-745".
- b. After the site Hospitals, add the qualifier as follows: Hospitals (non-occupied patient areas).
- c. On the front panel the statement should read: "For Household, Commercial/Agricultural Use."

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE
---------------------------------	------

- 2/4
- d. Change "Space Spray" to "Directed Spray" under instructions for Flea in Buildings. To this same paragraph, add "Allow all sprayed articles to dry thoroughly before use."
  - e. Change "Animal Use" to "Livestock Use" under Agricultural Uses.
  - f. There are no directions for treating pests thus delete the paragraph under Animals and add the following statement under Precautionary Statements:

Do not use on dogs or cats.

4. Submit five copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca  
Product Manager 13  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosure

# PYRAPERM

## INSECT KILLER WBA N67A

EPA REG. NO. 4816-745

EPA EST. NO. 279-NY-1

- \* Recommended for Household and Commercial/Agricultural Use
- \* Contains Pyrenone<sup>®</sup> and Permanone<sup>®</sup>
- \* A broad spectrum aqueous insecticide which kills ants, bed bugs, booklice, carpenter ants, carpet beetles, centipedes, cigarette beetles, clover mites, cluster flies, cockroaches, confused flour beetles, crickets, drug store beetles, fleas, flies, fruit flies, gnats, grain mites, granary weevils, horn flies, house flies, lice, millipedes, mosquitoes, mud daubers, red flour beetles, rice weevils, sawtoothed grain beetles, silverfish, small flying moths, sow bugs, spiders, stable flies, ticks and wasps.

### ACTIVE INGREDIENTS:

Pyrethrins .....	0.250%
*Piperonyl Butoxide, Technical .....	1.250%
**Permethrin .....	0.125%
INERT INGREDIENTS: .....	98.375%
	100.00%

\*Equivalent to min. 1.0% (butylcarbityl)(6-propylpiperonyl) ether and 0.25% related compounds.

\*\* (3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.

PYRAPERM, PYRENONE and PERMANONE - Registered Trademarks of Fairfield American Corporation.

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

See label for full directions.

NET CONTENTS: \_\_\_\_\_



FAIRFIELD  
AMERICAN  
CORPORATION

201 ROUTE 17 NORTH, RUTHERFORD, NJ 07070

745 022192A

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in and around Non Food/Feed Areas of (but not limited to) Beef Cattle Operations, Campgrounds, Catteries, Dairy Farms (including Milk House, Milk Parlor, Loading Sheds, and Holding Lot), Food Storage Areas, Hog Operations, Homes, Horse Barns and Stables, Hospitals, Hotels, Kennels, Motels, Poultry Operations, Resorts, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Utilities, Vehicles, Warehouses, and other Commercial and Industrial Buildings.

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held.

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Do not use in rooms/areas while occupied by patients, the elderly or infirm. Do not apply to classrooms when in use.

### SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches. Remove birds and cover fish aquariums before spraying.

### NONAGRICULTURAL USES

**Cluster Flies, Houseflies, Fruit Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Black Flies, Skipper Flies, Blow Flies and Small Flying Moths:** Close all doors and windows and spray at the rate of 1 to 3 seconds per 1000 cu. ft. moving toward the door. Fill room with mist. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Leave room and keep area closed for 15 minutes. Open and ventilate before reoccupying. Sweep up and destroy fallen insects. Repeat treatment as necessary.

**Ants, Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Clover Mites, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs and Pillbugs:** Spray into hiding places such as cracks and crevices, behind sinks, cabinets, meter boxes, door frames, windows, along baseboards and floors,

around drains and plumbing possible. Space spray: Open windows before spraying. Spray into all locations contacting as much space as possible and keep area closed for 15 minutes before reoccupying. Sweep up and destroy fallen insects.

**Indian Meal Moth, Chocolate Angoumois Grain Moth:** Spray into all parts of room and containers, pallets and dark areas for each 1000 cubic feet of space. Open and ventilate before reoccupying. Sweep up and destroy fallen insects. Repeat application as necessary.

**Exposed stages of Confused Flour Beetles, Drug Store Beetles, Flour Beetles, Rice Weevils:** Spray exposed adult and larval stages in crevices, processing and handling areas, containers or other areas with contact as many insects as possible. Contact 100 square feet of surface. Repeat application 1 hour after treatment. Space spray into all parts of room and keep area closed for 15 minutes before reoccupying. Repeat application weekly or as necessary. Dispose of, or treat with insecticide, infested articles to this purpose.

**Carpet Beetles:** Spray dishes, shelves and carpets. Repeat application as necessary.

**Bedbugs:** Spray mattresses, box springs, bed frames, moldings and floors. Repeat application as necessary. Remove sprayed articles to dry thoroughly.

To control **Clothes Moths** and **proteinaceous fibers:** Remove infested articles, brush thoroughly, and air for 24 hours. Apply spray liberally to empty storage areas directing the spray into all hiding places. Repeat application when insects are at rest. Contacting as many insects as possible.

**Mud Daubers and Wasp Nests:** Spray mud daubers and their nests from approximately 12 to 15 inches. Insects will fly away from fog. Apply spray when insects are at rest. Contacting as many insects as possible.

CONTINUED

dry

**Fleas in buildings (Space spray):** Hold can 36 inches above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq. ft. making sure that all floor, sofa and chair surfaces are contacted. Leave room and keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment after 7 days or as needed.

**Ticks (Adults and Larvae):** Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

**AGRICULTURAL USES:**

To control *Bees, Deer Flies, Face Flies, Fleas, Gnats, Horn Flies, Hornets, Horse Flies, House Flies, Lice, Mosquitoes, Small Flying Moths, Stable Flies, Ticks, and Wasps:*

**IN DAIRY FARMS**

**MILKING PARLOR AND MILK ROOM:** Cover milking utensils and milk to prevent contamination from spray and falling dead insects. Close windows and doors and direct fog upward in all directions at a rate of 1 to 3 seconds per 1000 cubic feet of space, moving toward the door. Leave room and for best results, keep area closed for 15 minutes after spraying. Do not remain in treated area. Open and ventilate after treatment is completed.

**ANIMAL USE (DO NOT USE ON SICK ANIMALS) -** From approximately 2 feet away, thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing. Apply for 2 to 4 seconds per animal. Repeat daily or as needed. To control *Blood Sucking Lice*, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

**IN STANCHION BARN**

Walk behind animals and direct spray over backs allowing 1 second per animal. Apply each morning for best results.

**IN BEEF CATTLE, GOAT, AND SHEEP OPERATIONS, HORSE BARN AND STABLES**

Close all windows and doors, and spray at a rate of 1 to 2 seconds per 1000 cu. ft. If area cannot be closed, double the dosage. Leave room and keep area closed for 15 minutes following application. Ventilate area after treatment is complete. **FOR ANIMAL APPLICATION (Do not use on sick animals),** from approximately 2 feet away, thoroughly spray entire animal, preferably as it is being released to pasture. Do not spray directly toward animal's face or eyes. Apply for 2 to 4 seconds per animal. Repeat daily or as needed. To control *Blood Sucking Lice*, apply to the

infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

**IN POULTRY OPERATIONS:**

Use only in unoccupied poultry houses when birds are not present. Close all windows and doors, and spray at a rate of 1 to 2 seconds per 1000 cu. ft, moving toward the door. If area cannot be closed, double the dosage. Leave room and keep area closed for 15 minutes following application. Ventilate area after treatment is complete. Treat daily or as needed.

To control *Poultry Lice*, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of *Bedbugs and Mites* in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

**IN HOG OPERATIONS:**

Close windows and doors and direct fog upward in all directions at a rate of 1 to 2 seconds per 1000 cubic feet of space, moving toward the door. If area cannot be closed, double the dosage. Leave room and for best results, keep area closed for 15 minutes after spraying. Do not remain in treated area. Open and ventilate after treatment is completed.

**HOG APPLICATION (DO NOT USE ON SICK ANIMALS) -** From approximately 2 feet away, thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's face or eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing. Apply for 1 to 2 seconds per hog. Repeat daily or as needed. Do not ship for slaughter within 5 days of last treatment. To control *Blood Sucking Lice*, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

**STORAGE AND DISPOSAL**

Store in a cool, dry area. Do not transport or store below 32°F.

Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**

Harmful if inhaled. Causes skin irritation. Do not get on skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish tanks before spraying. Vacate room after treatment

and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

**Animals:** Do not spray directly in/on eyes, mouth or genitalia of pets. Do not treat or cause exposure to kittens or puppies less than four weeks old.

**STATEMENT OF PRACTICAL TREATMENT**

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF INHALED:** Remove victim to fresh air. If not breathing, apply artificial respiration, preferably mouth to mouth. Get medical attention.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

**PHYSICAL OR CHEMICAL HAZARDS**

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not apply this product in conduits, motor housing, junctions and switch boxes or other electrical equipment because of possible shock hazard. Do not use on surfaces that can be harmed or stained by water.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

JUL 14 1992

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

4816-945

f/h