TETRAPERA

INDUSTRIAL CRAWLING INSECT KILLER N104

- * FOR COMMERCIAL / INDUSTRIAL USE
- * CONTROLS INDUSTRIAL AND GARDEN INSECTS
- FOR COMMERCIAL USE IN NON-FOOD/FEED AREAS OF INDUS-TRIAL AND INSTITUTIONAL BUILDINGS: SCHOOLS, WARE-HOUSES, OFFICE BUILDINGS, THEATERS, GARAGES, HOTELS, MOTELS.
- A BROAD SPECTRUM INSECTICIDE HIGHLY EFFECTIVE AGAINST ANTS, BEETLES (INCLUDING CARPET BEETLES, FLOUR BEETLES, GRAIN BEETLES AND GROUND BEETLES) CENTIPEDES, CLOVER MITES, COCKROACHES, CRICKETS (INCLUDING MOLE CRICK-ETS) EARWIGS, FIREBRATS, FLEAS, MILLIPEDES, MOTHS, PILL-BUGS, SCORPIONS, SILVERFISH, SOWBUGS, SPIDERS, TICKS AND MANY SIMILAR PESTS.
- * WATER-BASED FORMULA
- **NON-STAINING**

ACTIVE INGREDIENT:

*Tetramethrin	0.20%
**Permethrin	0.20%
***Piperonyl Butoxide, Technical	1.00%
TINERT INGREDIENTS:	
	100,00%

- *1-cyclohexene-1,2-dicarboximidomidomethyl-2,2-dimethyl-3-(2-met hylpropenyl)cyclopropanecarboxylate
- **(3-phenoxyphenyl)methyl(+/-)cis/trans 3-(2,2-dichloroethenyl)2,2-dimethylcyclopropanecarboxylate.

Cistrans ratio: min.35%(+/-)cis and max. 65%(+/-)trans.

*Equivalent to min. 0.80%(buylcarbityl)(6-propylpiperonyl)ether and 0.20% related compounds.

†Contains Petroleum Distillates


TETRAPERM - Trademark of AgrEvo Environmental Health, Inc.

EPA REG. NO. 4816-761

EPA EST. NO.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Side Panel For Use Cautions



CODE 82652

PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals

Avoid contact with skin, eyes and clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before space spraying or surface applications. Do not use in classrooms while in use. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

Environmental Hazards

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

Physical Or Chemical Hazards
Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not apply this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. Do not use on surfaces that can be harmed or stained by water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store in a cool, dry area. Do not transport or store below 32°F. Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

Do not use in food/feed areas of food handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held.

Nonfood/feed areas are areas such as garbage rooms, lavatories, floor_drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches.

INDOORS:

COCKROACHES, WATERBUGS, PALMETTO BUGS, SILVER-FISH, FIREBRATS, CRICKETS, SCORPIONS, SPIDERS, MILLI-PEDES, CENTIPEDES, SOWBUGS AND PILLBUGS, Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around grains and plumbing, hitting insects with spray whenever possible. Repeat treatment as nec-

ANTS: Spray trails, nests and points of entry. Spray on Ants where possible. Repeat as necessary.

TETRAPERMTM

INDIANMEAL MOTHS, ANGOUMOIS GRAIN MOTHS: Close doors and windows, direct spray into all parts of room especially around stored food/feed product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application as necessary.

Exposed stages of Confused Flour Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary.

CARPET BEETLES: Spray directly on *Beetles*, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

TO CONTROL FLEAS AND BROWN DOG TICKS: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor coverings. Fresh bedding should be placed in animal quarters following treatment. Do not spray pets with this product. Treat pets with a product registered for application to pets to control the source of flea infestation.

BEDBUGS: Spray mattresses, particularly around tuits and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary. Allow mattresses and other articles to dry thoroughly before use.

TO CONTROL CLOTHES MOTHS AND CARPET BEETLES on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be reated by spraying lightly, keeping nozzle at least three feet from tabric to avoid staining.

DUTDOORS:

Point spray nozzle away from face. Hold can on slight downward angle. For best results, spray when air is calm. Allow a few minutes for product to take effect. Spray with wind if breeze is blowing.

PESTS OUTSIDE SURFACE OF BUILDINGS: For control of building infestation due to Ants, Clover Mites, Crickets, Earwigs, Millipedes amd Sowbugs (Pillbugs). Spray foundation of building where insects are active and may find entrance, up to a height of 2 to 3 feet. Apply as a residual spray to outside surfaces of buildings where insects tend to congregate. Spray areas include but are not limited to) screens, window frames, eaves, porches, patios, garages and refuse dumps. Repeat treatment as necessary.

CRAWLING INSECTS: Ants, Cockroaches, including Asian Cockroaches, Crickets, Centipedes, and Silverfish. Spray infested surface of patio or picnic area, hitting as many insects as possible. Also spray legs of tables and chairs to repel Ants. Spray ant hills for best control.

ANTS, LONE STAR TICKS, DOG TICKS, CRICKETS, FLEAS AND EARWIGS: Apply thoroughly to infestation in bushes, grass or weeds. For *Ants*, spray freely around ant trails and hills.

FOR THE CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahiias, delphiniums, dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 °F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

AgrEvo Environmental Health 95 Chestnut Ridge Road Montvale, NJ 07645