### PYRAPERM® HOUSEHOLD INSECT KILLER WBA P61

- \* CONTAINS PYRENONE® AND PERMANONE®
- \* SEEKS OUT BUGS WHERE THEY HIDE
- \* KILLS ON CONTACT
- \* CONTROLS RESIDENTIAL AND GARDEN INSECTS
- \* FOR HOUSEHOLD USE
- \* NON-STAINING
- \* A BROAD SPECTRUM INSECTICIDE HIGHLY
  EFFECTIVE AGAINST ANTS, (INCLUDING
  CARPENTER ANTS), BEES, BEETLES(INCLUDING
  CARPET, FLOUR, GRAIN, AND GROUND
  BEETLES), CENTIPEDES, CLOVER MITES,
  COCKROACHES, CRICKETS (INCLUDING MOLE
  CRICKETS), EARWIGS, FIREBRATS, FLEAS,
  MILLIPEDES, MOTHS, PILLBUGS, SCORPIONS,
  SILVERFISH, SOWBUGS, SPIDERS, TICKS
  (INCLUDING TICKS WHICH MAY CAUSE LYME
  DISEASE), WASPS, AND OTHER PESTS LISTED
  ON THE LABEL
- \* KILLS LICE AND THEIR EGGS
- \* FOR USE ON BEDDING, FURNITURE AND OTHER INANIMATE OBJECTS INFESTED WITH LICE
- \* NO UNPLEASANT ODOR
- WATER-BASED FORMULA
- \* FOR USE ON DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS
- \* CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS
- \* PROTECTS FOR UP TO 4 WEEKS
- \* PROTECTS FOR UP TO 12 WEEKS
- \* PROTECTS FOR UP TO 3 MONTHS

#### **ACTIVE INGREDIENTS:**

Pyrethrins	0.20%
*Piperonyl Butoxide, Technical	
**Permethrin	0.40%
INERT INGREDIENTS:	98.40%
	100.00%

\*Equivalent to min. 0.8% (butylcarbityl)(6-propylpiperonyl) ether and 0.2% related compounds.

\*\*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2- dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min, 35% (+/-) cis and max, 65% (+/-) trans.

PYRAPERM, PYRENONE and PERMANONE - Registered Trademarks of AgrEvo Environmental Health, Inc.

EPA REG. NO. 4816-769

EPA EST. NO.

## KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT IE INHALED: Redrove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth, Get medical attention.

Under the Sede Re are Panel For Use Cautions
Fungicide, and households Act
as amended, for the pesticide
registered under
RPA Reg. No. 4816-769

AUG 2.9 1996

NET CONTENTS:

# PRECAUTIONARY STATEMENTS Hazards To Huggins & Domestic Animals CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash before reuse.

Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

#### **Environmental Hazards**

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

#### Physical Or Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Do not use in rooms/areas while occupied by patients, the elderly or infirm. Do not apply to classrooms when in use.

#### SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 in

#### INDOORS:

CLUSTER FLIES, HOUSEFLIES, FRUIT FLIES, MOSQUITOES, GNATS, WASPS, HORNETS, BEES, BLACK FLIES, SKIPPER FLIES, BLOW FLIES AND SMALL FLYING MOTHS: Close all doors and windows and direct spray upward into center of room with a slow sweeping motion. Spray 5-10 seconds for average room, moving toward the door. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Leave room and keep area closed for 15 minutes after spraying. Ventilate room therefughly before re-entry. Sweep up and discard faller insects. Repeat treatment as necessary.

COCKROACHES, WATERBUGS, PALMETTO BUGS, SILVERFISH, FIREBRATS, CRICKETS, SCORPIONS, SPIDERS, MILLIPEDES, CENTIPEDES, SOWBUGS, AND PILLBUGS: Spray into hiding places such as cracks and crevices, behind sinks, cabinots, along

baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary. Allow all sprayed articles to dry thoroughly before use.

TO CONTROL CLOTHES MOTHS AND CARPET BEETLES: on wool and other proteinaceous fibers: Remove infested articles from storage, brush choroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining

FLEAS IN BUILDINGS - SPACE SPRAY: Hold can 38 inches above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq. ft. making sure that all floor, sofa and chair surfaces are contacted. Leave room and keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment after 7 days or as needed.

FLEAS & TICKS, INCLUDING TICKS WHICH MAY CAUSE LYME DISEASE (ADULTS AND LARVAE): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture and mattresses may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

TO CONTROL FLEAS, TICKS AND LICE ON DOGS. Starting at the head, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur back to the tail. Repeat application weekly if necessary to control infestations.

NOTE: Not recommended for use on puppies less than four weeks old.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square

foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry moroughly before use.

IN ANIMAL QUARTERS: For rapid control of House Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees and Small-Flying Moths close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

TO CONTROL Blood Sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

#### **OUTDOORS:**

Point spray nozzle away from face. Hold can on slight downward angle. For best results, spray when air is calm. Allow a few minutes for product to take effect. Spray with wind if breeze is blowing.

FLYING INSECTS: Flies, Mosquitoes, Gnats, Small Flying Moths, Homets and Yellow Jackets. Allow fog to drift over area. Direct spray toward bushes, or grass from at least 4 to 6 feet away. Give special attention to breeding areas such as garbage cans and compost piles. Repeat as necessary.

MUD DAUBERS AND WASP NESTS: Spray mist directly on insects and their nests from approximately 2 feet away. Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Spray into hiding and breeding places contacting as may insects as possible.

CRAWLING INSECTS: Ants, Cockroaches, including Asian Cockroaches, Crickets, Mole Crickets, Centipedes, and Silverfish. Spray infested surface of patio or picnic area, hitting as many insects as possible. Also spray legs of tables and chairs to repel ants. Spray ant hills for best control.

FOR THE CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, camaticus, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds,

mododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

#### STORAGE AND DISPOSAL

Store in a cool, dry area. Do not transport or store below 32°F.

Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

AgrEvo Environmental Health 95 Chestnut Ridge Road Montvale, NJ 07645

)

- 3-7