


UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 4822-116

**WARNING: KEEP OUT OF REACH OF CHILDREN. EYE IRRITANT.**  
See first aid and additional precautions on back.

**Johnson**



**For Meat Cutting and  
Food Preparation Areas**

U

**ONE GALLON**

A PRODUCT OF **Johnson wax** SERVICE PRODUCTS DIVISION

ACTIVE INGREDIENTS: n-Alkyl (60% C<sub>14</sub>, 30% C<sub>16</sub>, 5% C<sub>12</sub>, 5% C<sub>18</sub>) dimethyl  
benzyl ammonium chlorides 1.29%; n-Alkyl (68% C<sub>12</sub>, 32% C<sub>14</sub>) dimethyl ethyl-  
benzyl ammonium chlorides 1.29%.

INERT INGREDIENTS: 97.42%.

445658

ED  
TICIDE  
E ACT  
ISTER-

# Johnson J-80 sanitizer

Johnson J-80 Sanitizer is expressly formulated for the effective sanitizing of food processing equipment and work areas which have been previously cleaned. It is recommended for use in all areas where meat cutting, food preparation and other food services take place in stores and in commercial restaurants and kitchens.

J-80 saves time and labor because no after-rinse is required when used at the recommended concentration of 200 ppm of active quaternary. J-80 may be applied with either mechanical or manual spraying equipment. J-80 is effective as a sanitizer when diluted in water of up to 700 ppm hardness (as CaCO<sub>3</sub>).

## DIRECTIONS FOR USE

**For sanitizing surfaces of food processing equipment and work areas.**

1. Clean all surfaces thoroughly with Johnson Breakup Cleaner. Tilt all movable surfaces for proper drainage. Rinse with hot water.
2. Mix 1-oz. J-80 Sanitizer with each gallon of water.
3. Wet surfaces thoroughly with J-80 solution.
4. Let surfaces drain and air dry. Do not rinse. Do not wipe.

**For sanitizing multi-use eating and drinking utensils**

1. Scrape and pre-wash utensils and glassware whenever possible.

2. Wash with a good detergent.
3. Rinse with clean water.
4. Sanitize in a solution of 1 oz. of J-80 to each gallon of water (200 ppm). Immerse all utensils for at least 2 minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drain board to air dry. Do not rinse. Do not wipe.


**NOTE:** Change sanitizer solution daily. Periodically test sanitizer solution for a concentration of 200 ppm active quaternary by using the test kit available for this purpose.

**WARNING:** Keep out of reach of children. Causes eye or skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food. Rinse empty container thoroughly with water and discard it. Avoid inhalation of spray mist.

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of water. Avoid alcohol. Call a physician immediately.

© S. C. JOHNSON & SON, INC., RACINE, WIS. 53403, U.S.A.

 Johnson and J-80 are trademarks

A PRODUCT OF  SERVICE PRODUCTS DIVISION

EPA Est. 4822-WI-1

EPA Reg. No. 4822-116-AA

**KEEP FROM FREEZING**

445659