

P11/ 31 4822-358 1948  
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

JUL 24 1995

S.C. Johnson & Son, Inc.  
1525 Howe Street  
Racine, WI 53403

Attention: Stephen Smith  
Senior Registration Specialist

Subject: Virex II Ready To Use (RTU)  
EPA Registration No. 4822-358  
MOU Compliance Amendment dated June 12, 1995

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, will be acceptable provided that you make the labeling change listed below before you release the product for shipment bearing the amended label.

1. Under the "Directions for Use", item number 2, and the statement, "Allow the surfaces to remain wet for ten minutes."

A stamped copy is enclosed for your records. Submit five (5) copies of the final printed label prior to release of the product for shipment.

If you have any questions concerning this letter, please contact Wanda Mitchell at (703) 305-6141.

Sincerely yours,



Marion J. Johnson, Jr.  
Product Manager (31)  
Antimicrobial Program Branch  
Registration Division (7505C)

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

2048

VIREX II READY TO USE (RTU)  
One Step Germicidal Cleaner and Deodorant

Pseudomonacidal Staphylocidal Salmonellacidal  
Fungicidal \*Virucidal

ACTIVE INGREDIENTS:

Didecyl dimethyl ammonium chloride . . . . .	0.034%
n-Alkyl [C <sub>14</sub> 50%, C <sub>12</sub> 40%, C <sub>16</sub> 10%] dimethyl benzyl ammonium chloride . . . . .	0.032%
INERT INGREDIENTS . . . . .	99.934%

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

See additional precautionary statements on back

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

JUL 24 1995

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No.

4322-358

VIREX II Ready To Use (RTU) is a germicidal cleaner and deodorant that requires no dilution before use (ready to use). VIREX II Ready To Use (RTU) is labor saving, providing broad spectrum disinfection, mold and mildew control, and odor elimination with mild cleaning.

VIREX II Ready To Use (RTU) is recommended for use on non-food contact surfaces in hospitals, nursing homes, schools and colleges, veterinary clinics, animal life science laboratories, hotels and federally inspected meat and poultry establishments.

VIREX II Ready To Use (RTU) is specially formulated to be non-dulling to floor finishes. VIREX II Ready To Use (RTU) eliminates the time and labor normally required for rinsing.

VIREX II Ready To Use (RTU) is bactericidal against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella choleraesuis*, *Staphylococcus aureus* (methicillin resistant - MRSA), *Escherichia coli*, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Proteus mirabilis*, *Proteus vulgaris*, *Salmonella enteritidis*, *Salmonella pullorum*, *Salmonella typhi*, *Serratia marcescens*, *Streptococcus agalactiae*, *Streptococcus faecalis*, and *Streptococcus pyogenes* according to the AOAC Use Dilution Test Method modified in the presence of 400 ppm synthetic hard water (calculated as  $\text{CaCO}_3$ ) plus 5% organic serum.

VIREX II Ready To Use (RTU) is fungicidal against *Trichophyton mentagrophytes*, *Candida albicans* and *Aspergillus niger* according to the AOAC Use Dilution Test, modified in the presence of 400 ppm hard water (calculated as  $\text{CaCO}_3$ ) plus 5% organic serum.

VIREX II Ready To Use (RTU) is virucidal against \*HIV-1 (AIDS virus), Influenza A2/J305, Herpes Simplex Type 1, Herpes Simplex Type 2, Adenovirus Type 2, New Castle disease virus, Avian influenza, and Pseudorabies virus according to the virucidal qualification test, modified in the presence of 400 ppm hard water (calculated as  $\text{CaCO}_3$ ) plus 5% organic serum.

#### DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

#### GENERAL USE DIRECTIONS FOR DISINFECTING:

For use on hard inanimate non-porous surfaces such as floors, walls, metal surfaces, porcelain, and plastic surfaces. Remove gross filth and heavy soil deposits then thoroughly wet surfaces as required. Use full strength (no dilution) for a minimum contact time of 10 minutes in a single application. Can be applied with a mop, sponge, or cloth as well as coarse spraying or soaking. Rinsing is not necessary unless floors are to be waxed or polished. Use a fresh solution daily or when use solution becomes visibly dirty.

\* Kills HIV-1 (AIDS virus) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS virus) ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

4078  
PERSONAL PROTECTION: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

JUL 24 1995

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

4822-358

5048

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

CONTACT TIME: HIV-1 (AIDS virus) is inactivated after one (1) minute contact time.

Use a ten (10) minute contact time for disinfection against all other bacteria, fungi, and viruses listed on labeling.

DISPOSAL OF INFECTIOUS MATERIAL: Blood and other body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

##### CAUTION

Causes eye irritation. Avoid contact with skin, eyes, and clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling.

#### STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

#### PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

#### CONTAINER DISPOSAL

##### Plastic Containers

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

##### Fiber Drums with Liners

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

##### Metal Containers

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

JUL 21 1995

(If container is 1 gallon or less, use this container disposal statement.)  
Do not reuse empty container. Wrap container and put in trash.

6078

General

Consult Federal, State, or local disposal authorities for approval of alternative procedures such as limited open burning.

EPA Registration No. 4822-358  
EPA Est. No. 4822-WI-1

Net Contents 32 oz. - 55 gallons

1992 S. C. Johnson & Son, Inc. Racine, WI 53403-5011, U.S.A.  
All Rights Reserved

JUL 24 1995

4822-358

REFERENCE SHEET

VIREX II READY TO USE (RTU)

ONE STEP GERMICIDAL CLEANER AND DEODORANT

VIREX II Ready To Use (RTU) is a germicidal cleaner and deodorant that requires no dilution before use (ready to use). VIREX II Ready To Use (RTU) is labor saving, providing broad spectrum disinfection, mold and mildew control, and odor elimination with mild cleaning.

Effective against the following pathogenic bacteria according to the AOAC Use Dilution Test, 15th edition, 1990, modified in the presence of 400 ppm synthetic hard water (calculated as  $\text{CaCO}_3$ ) plus 5% organic serum:

- |  |   |
|--|---|
| Pseudomonas aeruginosa ATCC# 15442           | <sup>1</sup> Salmonella pullorum PI# 3751     |
| Salmonella choleraesuis ATCC# 10708          | Salmonella typhi ATCC# 6539                   |
| Staphylococcus aureus ATCC# 6538             | Serratia marcescens ATCC# 14756               |
| <sup>4</sup> Escherichia coli 0157:H7        | <sup>3</sup> Staphylococcus aureus SCJ# 226   |
| Klebsiella pneumoniae ATCC# 4352             | (methicillin resistant - MRSA)                |
| <sup>4</sup> Listeria monocytogenes 20A OHIO | <sup>2</sup> Staphylococcus aureus SCJ# 225   |
| <sup>1</sup> Proteus mirabilis PI# 366       | <sup>1</sup> Streptococcus agalactiae PI# 517 |
| Proteus vulgaris ATCC# 13315                 | <sup>1</sup> Streptococcus faecalis PI# 523   |
| Salmonella enteritidis ATCC# 13076           | Streptococcus pyogenes ATCC# 9342             |
| <sup>4</sup> Salmonella enteritidis E40      |   |

Fungicidal against Trichophyton metagrophytes ATCC# 9533, Candida albicans ATCC# 10231, and Aspergillus niger ATCC# 6275 according to the AOAC Use Dilution test, 15th edition, 1990, modified in the presence of 400 ppm hard water (calculated as  $\text{CaCO}_3$ ) plus 5% organic serum.

Virucidal against \*HIV-1 (AIDS virus), Adenovirus Type 2 ATCC# VR-2, Herpes Simplex Type 1 ATCC# VR-735, Herpes Simplex Type 2 ATCC# VR-734, Influenza A2/J305 ATCC# VR-100, New Castle disease virus ATCC# VR623, Avian influenza ATCC# VR797 and Pseudorabies virus ATCC# VR135 according to the Virucidal Qualification test, modified in the presence of 400 ppm hard water (calculated as  $\text{CaCO}_3$ ) plus 5% organic serum.

Note: Clinical isolates of organisms designated by footnotes.

- <sup>1</sup> Presque Isle Associates
- <sup>2</sup> St. Mary's Hospital, Racine, Wisconsin
- <sup>3</sup> Park West Hospital, Knoxville, Tennessee
- <sup>4</sup> Food Research Institute, Madison, Wisconsin

W:  
in EPA ..

JUL 24 1995

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended for the pesticide  
review under EPA Reg. No.

482-358

8078

\* Kills HIV-1 (AIDS virus) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS virus) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

**CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

**CONTACT TIME:** HIV-1 (AIDS virus) is inactivated after one (1) minute contact time.

1 Use a ten (10) minute contact time for disinfection against all other bacteria, fungi, and viruses listed on labeling.

**DISPOSAL OF INFECTIOUS MATERIAL:** Blood and other body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No  
4822-358