


FOR INDUSTRIAL USE ONLY

IMPROVED NELCOPHENE

DANGER

EPA
Registration Number 4823-6



Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food. ct,

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

Cameraman's Note
Poor Copy

NOTE TO

PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

TUBERCULOCIDAL	GERMICIDAL	DEODORANT	PSEUDOMONOCIDAL
Recommended for Conductive Floors			
MINIMUM Phenol Coefficient (AOAC) S. Typhosa (20°C) 37		S. Aureus (20°C) 32	
Minimum Use Dilution: Confirmation (AOAC) S. Choleraesuis 1:450		S. Aureus 1:450 Ps. aeruginosa 1:128	
Active Ingredients			
	Isopropyl Alcohol		10.00%
	Potassium Laurate		10.00%
	Sodium o-benzyl-p-chlorophenate		6.60%
	Sodium p-tertiary-amylphenate		5.67%
	Sodium o-phenylphenate		3.30%
	Tetrasodium ethylenediamine tetraacetate		1.50%
	Sodium salt of 2,2-Methylene Bis (3, 4, 6- trichlorophenol)		5.56%
			37.72%
Inert Ingredients			
			62.28%
			100.00%

IMPROVED NELCOPHENE is a broad spectrum, general disinfectant. It may be used in hospitals, operating rooms, nursing homes or medical institutions. It may be used in tuberculocidal areas. Improved Nelcophene is also recommended for use in homes, hotels, motels, and schools as well as in food processing and packing establishments.

When used according to instructions Improved Nelcophene is not harmful to washable, painted surfaces, resilient flooring, metal surfaces or terrazzo.

RECOMMENDED DILUTIONS AND METHOD OF APPLICATION:
When used in hospitals, operating rooms, nursing homes or food establishments heavy deposits of grime or grease should be

PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

TUBERCULOCIDAL	GERMICIDAL	DEODORANT	PSEUDOMONOCIDAL
Recommended for Conductive Floors			
MINIMUM Phenol Coefficient (AOAC) S. Typhosa (20°C) 37		S. Aureus (20°C) 32	
Minimum Use Dilution Confirmation (AOAC) S. Choleraesuis 1:450		S. Aureus 1:450 Ps. aeruginosa 1:128	
Active Ingredients			
	Isopropyl Alcohol		10.00%
	Potassium Laurate		10.00%
	Sodium o-benzyl-p-chlorophenate		6.60%
	Sodium p-tertiary-amylphenate		5.67%
	Sodium o-phenylphenate		3.39%
	Tetrasodium ethylenediamine tetraacetate		1.50%
	Sodium salt of 2,2-Methylene Bis (3, 4, 6- trichlorophenol)		0.56%
			37.72%
Inert Ingredients			
			62.28%
			100.00%

Cameraman's Note

Poor Copy

IMPROVED NELCOPHENE is a broad spectrum, general disinfectant. It may be used in hospitals, operating rooms, nursing homes or medical institutions. It may be used in tuberculocidal areas. Improved Nelcophene is also recommended for use in homes, hotels, motels, and schools as well as in food processing and packing establishments.

When used according to instructions Improved Nelcophene is not harmful to washable, painted surfaces, resilient flooring, metal surfaces or terrazzo.

RECOMMENDED DILUTIONS AND METHOD OF APPLICATION:

When used in hospitals, operating rooms, nursing homes or food establishments heavy deposits of grime or grease should be removed with a mild detergent. Using one ounce of Improved Nelcophene per gallon of clean water thoroughly mop or scrub surfaces. This concentration is effective against Ps. aeruginosa.

When Improved Nelcophene is used to disinfect food contact surfaces they should be thoroughly scrubbed with a detergent and rinsed with potable water prior to reuse.

For tuberculocidal areas using same method of application use 2/3rds of a fluid ounce per gallon of water.

When used to store heat sterilized surgical instruments use 1/4 fluid ounces of Improved Nelcophene per gallon of water.

Minimum Use Dilution Confirmation against T. Interdigitale is 1:200 in 5 minutes.

Rinse empty container thoroughly with water and discard it.

TESTED
OCT 21 1977
FEDERAL INSECTICIDE
AND FUNGICIDE ACT
EDISON REGISTER-
4823-6 SUBJECT
10 ATTACHED COMMENTS

Manufactured by
MAINTENANCE SUPPLY SERVICE CORP.
HUNTERSVILLE, NORTH CAROLINA

Net Contents gal.