FORMULA #96

NET CONTENTS

GALLONS

WEED KILLER

Contains 1/4 pound of Urox per Gallon
For General Weed Control on Non-Crop Lands and Paved Areas

USDA Reg. No. 4823-13

ACTIVE INGREDIENTS:

Monuron Trichleroacetate
[3-(p-chlorophenyl)-1, 1-dimethylurea trichleroacetate] 3.19%

Aromatic Petroleum Derivative 91.67%

INERT INGREDIENTS 5.14%

CAUTION — Keep out of reach of children.

For Additional Caution Information See Right Panel
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

MADE IN U.S.A.

PRINTED IN U. S. A

Manufectured by

MAINTENANCE SUPPLY SERVICE CORP.

HUNTERSYILLE, NORTH CAROLINA

Dec-8-1970

4833-13-

DIRECTIONS

NON-CROP AREAS

For use on Non-crop land such as Right-of-Ways, Industrial Plants, Lumber Yards, Tank Farms, Storage Areas, and Parking Lots for control of most annual and perennial grasses and broadleaved weeds. The effectiveness and length of time that control will last will depend on type of soil, amount of rain and other conditions.

RATE: Under ordinary conditions use at the rate of 2 to 4 gallons per 1000 square feet (1-3/5 to 3-1/5 pints per 100 square feet). Under conditions where the weeds are difficult to control, use at the rate of 6 gallons per 100 square feet. (4-4/5 pints per 100 square feet).

HOW TO APPLY: Use a sprayer which will distribute the required amount evenly over the area to be treated. Use a coarse spray to reduce drift. Application may be made when top growth is dense, but better control will be obtained when weeds are short or when excess growth is mowed and removed before applying.

WHEN TO APPLY: Best results will be obtained if applied shortly before or at the time plant growth begins.

Rain or thorough watering after application will cause the material to move to the root zone and thus speed up control.

NOTE: Do Not apply, allow to drift or drain or flush equipment onto or near croplands, lawns, trees or other desirable plants or to areas where their roots may extend. Do Not contaminate irrigation ditches or water used for domestic purposes. Do Not allow contact with fertilizers, insecticides, fungicides or seeds. Application equipment should be thoroughly cleaned with a detergent and water to prevent corrosion. Equipment which the been used to apply this product should not be used for applying insecticides, fungicides, fertilizers or other agricultural chemicals as plant injury may result.

DESTROY CONTAINER WHEN EMPTY

Flush out thoroughly with water and drain where flush water can be run off without harm. Destroy container so that it may not be used again for any purpose and dispose of it at some location where no attempt to salvage will be made.

CAUTION:

HARMFUL IF SWALLOWED IRRITATING TO SKIN AND EYES

Do not get on skin, in eyes or on clothing.

In case of contact, immediately wash skin with plenty of water: for eyes, wash with water for at least 15 minutes and get medical attention.

Follow directions carefully. Timing and method of application is there indicates conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not to strict accordance with directions given herewith.

FORMULA #96 BEST DOCUMENT AVAILABLE

NET CONTENTS GALLONS

WEED KILLER

Contains 1/4 pound of Urox per Gallon For General Weed Control on Non Crop Lands and Paved Areas USDA Rep. No. 482 (13

ACTIVE INGREDIENTS:

Monuron Problems tate

[3-ip-chturoph-sy: 1 1 2 a ethylures trichlores, but 12 a Aromatic Fetre & historication 9167 INERT INGREDIENTS

CAUTION - Keep out of reach of children.

For Additional Caution Information See Right Paget DO NOT USE OR STORE NEAR HEAD OR GEEN FLAMI

MADE IN USA

5- 3-63 4 6 ...

Menufactured by MAINTENANCE SUPPLY SERVICE CORP.

> Acres 10 Direct 1971

1.32.72-

DIRECTIONS

NON-CROP AREAS

For use on Non-crop tand such as Righted-Ways, Industrial Plants, Lumber Yards, Tank Farms, Shares dreas, and Parking Lots for central of most annual and perennial grasses and broadlessed wheels. The effectiveness and length of time that central will last mill depend on type of soil, amount of rain and other conditions.

RATE If we introduce the condition is so at the rate of 2 to 6 gallons per 1000 square feet (1-3/5 to 3-1/5 pints per 100 square feet will 5 pints per 10 is square feet). How to apply Use a sprayer which will distribute the required amount eventy over the area to be treated. Use a course pray to reduce drift Application may be made when top growth is dense, but better control will be obtained when weeds are short or when excess prowth is moved and removed before applying.

WHEN TO APPLY Real regulate with the released of across prowth before or at the time plant growth begins.

WHEN TO APPLY: Best results with the obtained of applied shortly before or at the time plant growth begins.

Rawn or thorough watering after application will cause the material to move to the root zone and thus speed up control.

NOTE: Do Not apply, a particle of the rotation of lack continent entology near croplands, lawns, trees or other describle plant or to areas where their interests and not contaminate irrigation ditches or water used for demestic purposes. Do Not allow instant with the trees in entirides fungicides or seeds. Application equipment should be thoroughly clear a with a contamination which the property and water to prevent corrosion. Equipment which the property of the property of

BESTROY CONTAINER WHEN EMPTY

From our function with water and the water was to be un off without harm. Destroy container so that it may not be a formation where no attempt to

CAUTION!

HARMFUL IF SWALLOWED IRRITATING TO SKIN AND EYES Do not get on skin, in eyes or on clothing

In case of contact, immediately was I skin with plenty of water: for eyes, wash with water for at least 15 minutes and get medical attention.

for the product are beyond the control of the contr

