

ACCEPTED
 Jul 12, 1971
 4328-37

W. K. - DT133

DIRECTIONS FOR USE

NOTE: Keep spray tank agitated.

W. K. - DT 133 controls following weeds:

Aspens, Blackberries, Chokecherry, Cottonwood, Coyote Brush, Currants, Elm, Elderberry, Honeysuckle, Horsenettle, Locust, Maples, Milkvetch, Oaks, Plums, Poison Ivy, Perennial Groundcherry, Raspberries & related brambles, Sassafras, Sumac, Sweetgum, Wild Gooseberries, Wild Roses, Willows, Wild Grapes and many other species.

CONTROL OF WOODY PLANTS UP TO 6 TO 8 FEET TALL -- Mix this product at the rate of 3 to 4 quarts per 100 gals. of water. Spray when in full leaf, preferably in early summer. Wet plants thoroughly. Complete control usually requires spot-treating of survivors and seedlings during successive years. When present, grasses quickly replace brush and weeds.

CONTROL OF WOODY PLANTS OVER 8 FEET TALL -- Mix this product at the rate of 4 to 8 gals. per 100 gals. of oil. Apply preferably by Basal Bark Spray at any season of the year, but the dormant season is preferable for selective killing. Spray basal 12 to 20" around each stem or trunk, also exposed roots and crown area. Wet completely to point of run-off. Treated trees and shrubs die rather slowly, and should not be cut for a full season after treatment. Hard to kill woody plants may require repeat applications for effective control if new growth appears. Vapors from this product may injure susceptible plants in the immediate vicinity.

TREATMENT OF STUMPS: Mix this product at rate of 4 to 8 gals. per 100 gals. of oil. Spray to thoroughly wet tops, sides and exposed roots of freshly cut stumps to prevent sprouting. Trees with stumpage value and trees which will have to be removed can be treated in this fashion.

CAUTIONS

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Flush eyes with clear water. Call physician immediately. Avoid inhaling spray mist.

Avoid spray drift to susceptible plants, flowers, fruits, vegetables and ornamentals. Avoid contact with St. Augustine grass, Bent grass and clover due to possible injury to these fine grasses.

Use a coarse spray which is less likely to drift than a fine mist spray. Do not contaminate water used for irrigation or domestic purposes. Do not transport with or store near seeds, fertilizers, fungicides or insecticides. Do not use spray equipment for any other purpose, since traces of this product may be harmful to sprayed vegetation.

Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

Do not use around the home, recreation areas, or similar areas.

This product is toxic to fish. Keep out of lakes, streams or ponds, and do not contaminate water by cleaning of equipment or by disposal of wastes.

Do not reuse shipping containers; destroy by crushing and burying where possible. Store above 32°F. If frozen warm up to above 40°F. and agitate before using. Do not store with foodstuffs.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

ABCO ABCO ABCO



U. S. D. A. REG.
 NO. 4828-37

MANUFACTURING CHEMISTS
W. K. - DT 133

For control of certain woody plants and broad-leaved weeds in established grass pastures, rangelands, roadsides, rights-of-way, and other similar areas where desirable plants will not be injured.

ACTIVE INGREDIENTS:

Isooctylester of 2, 4-dichlorophenoxyacetic Acid 34.0%

Isooctylester of 2, 4, 5-trichlorophenoxyacetic Acid 32.5%

INERT INGREDIENTS: 33.5%

(Acid Equivalent: 2, 4-dichlorophenoxyacetic Acid 22.50% - 2 lb./gal.

2, 4, 5-trichlorophenoxyacetic Acid 22.50% - 2 lb./gal.

CAUTION

Keep Out of Reach of Children

(See direction label for other handling cautions)

NET CONTENTS GAL

Manufactured by

ABCO INC. P. O. BOX J IRWIN, PA. 15642 U.S.A.

REVISED 671