

July 12, 1971

4828-37

# W. K. - DT133

## DIRECTIONS FOR USE

W. K. - DT 133 controls following weeds:

Aspens, Blackberries, Chokecherry, Cottonwood, Coyote Brush, Currants, Elm, Elderberry, Honeysuckle, Horsechill, Locust, Maples, Milkweeds, Oaks, Plums, Poison Ivy, Perennial Groundcherry, Raspberries & related brambles, Sassafras, Sumac, Sweet gum, Wild Gooseberries, Wild Roses, Willows, Wild Grapes and many other species.

**CONTROL OF WOODY PLANTS UP TO 6 TO 8 FEET TALL** - Mix this product at the rate of 3 to 4 quarts per 100 gals. of water. Spray when in full leaf, preferably in early summer. Wet plants thoroughly. Complete control usually requires spot treating of survivors and seedlings during successive years. When present, grasses quickly replace brush and weeds.

**CONTROL OF WOODY PLANTS OVER 8 FEET TALL** - Mix this product at the rate of 4 to 8 gals. per 100 gals. of oil. Apply preferably by Basal Bark Spray at any season of the year, but the dormant season is preferable for selective killing. Spray basal 12 to 20" around each stem or trunk, also exposed roots and crown area. Wet completely to point of run-off. Treated trees and shrubs die rather slowly, and should not be cut for a full season after treatment. Hard to kill woody plants may require repeat applications for effective control if new growth appears. Vapors from this product may injure susceptible plants in the immediate vicinity.

**TREATMENT OF STUMPS:** Mix this product at rate of 4 to 8 gals. per 100 gals. of oil. Spray to thoroughly wet tops, sides and exposed roots of freshly cut stumps to prevent sprouting. Trees with stumpage value and trees which will have to be removed can be treated in this fashion.

NOTE: Keep spray tank agitated.

## CAUTIONS

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Flush eyes with clear water. Call physician immediately. Avoid inhaling spray mist.

Avoid spray drift to susceptible plants, flowers, fruits, vegetables and ornamentals. Avoid contact with St. Augustine grass, Bent grass and clover due to possible injury to these fine grasses.

Use a coarse spray which is less likely to drift than a fine mist spray. Do not contaminate water used for irrigation or domestic purposes. Do not transport with or store near seeds, fertilizers, fungicides or insecticides. Do not use spray equipment for any other purpose, since traces of this product may be harmful to sprayed vegetation.

Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.


Do not use around the home, recreation areas, or similar areas.

This product is toxic to fish. Keep out of lakes, streams or ponds, and do not contaminate water by cleaning of equipment or by disposal of wastes.

Do not reuse shipping containers; destroy by crushing and burying where possible. Store above 32 F. If frozen warm up to above 40 F. and agitate before using. Do not store with foodstuffs.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instruction.

ABCO ABCO ABCO



U. S. D. A. REG.  
NO. 4828-37

### MANUFACTURING CHEMISTS W. K. - DT 133

For control of certain woody plants and broad-leaved weeds in established grass pastures, rangelands, roadsides, rights-of-way, and other similar areas where desirable plants will not be injured.

**ACTIVE INGREDIENTS:**  
Isooctylester of 2, 4-dichlorophenoxyacetic Acid 34.0%  
Isooctylester of 2, 4, 5-trichlorophenoxyacetic Acid 32.5%  
**INERT INGREDIENTS 33.5%**  
(Acid Equivalent: 2, 4-dichlorophenoxyacetic Acid 22.50% - 3 lb./gal.  
2, 4, 5-trichlorophenoxyacetic Acid 22.50% - 3 lb./gal.)

**CAUTION**  
Keep Out of Reach of Children  
(See direction label for other handling cautions)

NET CONTENTS GAL

Manufactured by

ABCO INC. P. O. BOX J IRWIN, PA. 15642 U.S.A.

REVISED 6-71

**ACCEPTED**

7-12-71


UNDER THE FEDERAL INSECTICIDE,  
FUNGICIDE AND MITTICIDE ACT  
FOR CONTROL OF WEEDS AND  
PLANTS

**ABCO** ~~ABCO~~ ~~ABCO~~

4828-34

USDA REG.

NO. 4828-34



**MANUFACTURING CHEMISTS  
IMPROVED W.K-245**

For control of certain trees, noxious  
weeds, woody plants and brush in  
right-of-ways, pastures, rangeland,  
fence areas & similar non-crop areas.

**Active Ingredients:**

Isooctyl Ester of 2, 4, 5-Trichlorophenoxyacetic  
Acid\* ..... 64.2%

Inert Ingredients ..... 35.8%

\*Equivalent to 44.62% by weight of 2, 4, 5-Trichloro-  
phenoxyacetic acid, or not less than 4 lbs. of 2, 4,  
5-Trichlorophenoxyacetic acid per gallon at 68°F.

**CAUTION**

**Harmful if Swallowed**

**Keep Out of Reach of Children**

(See back panel for additional cautions)

CONTENTS

GALS.

This pro  
ing speci  
erous to  
2, 4, 5-T  
ated bra  
senettle,  
oak, Osa  
Perennia  
Waxmyr  
CONTR

Wet lea  
mixture  
100 gall  
are activ  
herbaceo  
be mad  
bloom. I  
and first  
4 feet in

CONTR  
W.K-245  
wet foli  
when in  
rapid gr  
fumes a  
year if

CONTR  
applicat  
W.K-245  
oil per  
during h  
to 80 d  
applying  
ed plan  
100 gal.  
age per  
peat ap  
mended  
ment to

BASAL  
MENTS  
gal Fue  
20 inch  
of run-c  
first, th  
stumps  
off to p  
trees. A

FRILL  
to kill  
Make si