


ABCO ABCO ABCO



USDA Reg. No. 4828-55

MANUFACTURING CHEMISTS

RODEX

KILLS RATS AND MICE

Active Ingredient:

Warfarin 2--(a-acetylbenzyl)---4---

Hydroxycoumarin 0.5%


Inert Ingredients: 99.5%

Total 100.0%

CAUTION

Keep Out of REACH of CHILDREN

NET CONTENTS POUNDS



RODEX

Kills Rats and Mice

FOR RATS: Mix thoroughly with 19 parts of any one of the following materials: (1) Corn meal, (2) Rolled Oats, (3) Dog Food, (4) Stock food, etc. Up to 5% of oil, such as mineral, corn, peanut or other edible oil may be used when the bait is so dry that it might tend to dust. If the bait material is made up with corn meal, rolled oats or oatmeal, a semi permanent bait may be prepared and need only be replaced when eaten by the rats. When you use perishable baits such as ground meat, fish, vegetables or similar material, place bait in position in late afternoon and replace daily with fresh bait. Dogs and cats will actively search for fish or meat baits, so place baits carefully to prevent feeding by these animals.

Place from 4 oz. to 1 lb. in areas frequented by rats or where rats have been known to feed. Replace baits as eaten. Maintain an uninterrupted bait supply for at least 10 days: Replace contaminated or spoiled baits promptly.

FOR MICE: Follow similar directions to above, but reduce amounts of bait placements to 1/2 oz., placed 8-12 foot intervals. Supply bait for at least 15 days. Continue baiting as long as there is evidence of feeding.

NOTE: Baiting should be continued as long as there is evidence that the rats or mice are still feeding on the bait. Under ordinary conditions this should take from 10 to 15

days. If there is a continuous source of infestation nearby, permanent bait stations should be utilized, and bait placed in them as regularly as may be necessary for conditions.

CAUTION

RODEX contains as its active ingredient an anti-coagulant chemical. Therefore, baits prepared using Rodex must be protected from or placed where humans, livestock, poultry or pets will not eat them. Do not use foods which will be attractive to children, such as cookies, candies, nuts, etc. If baits are eaten, give a tablespoon of salt in a glass of warm water and repeat until vomit is clear. Call a physician at once.

NOTE TO PHYSICIANS

When a human being has been known to ingest any RODEX, blood transfusions combined with intravenous injections and oral doses of Vitamin K should be prescribed or given to the patient as is usually prescribed in the instance of a hemorrhage caused by overdoses of Dicumarol.

Seller make no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of the material when such use and or handling is contrary to label instructions.

MANUFACTURED BY

ABCO, INC. P. O. BOX J IRWIN, PA. 15642 U. S. A.