


ABCO ABCO ABCO



USDA REG.
No. 4828-63

MANUFACTURING CHEMISTS

WKS - 65 BRUSH KILLER

ACTIVE INGREDIENTS:

(1) Isooctyl ester of 2,4 dichlorophenoxyacetic acid	46.0%
(2) Isooctyl ester of 2,4,5-Trichlorophenoxyacetic acid	10.5%
(3) Isooctyl ester of Silvex (2-(2,4,5-Trichlorophenoxy propionic acid	10.6%
INERT INGREDIENTS	32.9%
Total	100.00%

Equivalent to 30.79% by wt. or not less than 2-2/3 lbs. of 2,4-Dichlorophenoxyacetic acid per gallon; 7.43% by wt. or not less than 2/3 lbs. of 2,4,5-Trichlorophenoxyacetic acid per gallon; 7.66% by wt. or not less than 2/3 lbs. of Silvex per gallon.

CAUTION
Keep Out of Reach of Children
(Refer to Cautions on Back Panel)

CONTENTS _____ GALS.

WKS - 65

For control of woody plants, noxious weeds and broadleaf weeds along roadsides, rights of way, fence lines, under utility lines, vacant lots and similar areas where desirable plants and grass will not be damaged.

CAUTION
Keep Out of Reach of Children

HARMFUL IF SWALLOWED
Avoid contact with skin, eyes or clothing. Do not take internally. In case of contact flush eyes with plenty of water; wash skin with soap and water. Avoid breathing spray mist. Do not allow solution or spray drift to contact desirable plants, trees or ornamentals. Coarse sprays are less likely to drift. It is difficult to completely clean WKS-65 from sprayers or other equipment, therefore, do not reuse equipment for purposes where even trace amounts of this chemical may cause injury. Do not reuse shipping containers; destroy by crushing and burying. Avoid contamination of water intended for irrigation and domestic use. Do not transport with or store near seeds, fertilizers, insecticides or fungicides. Vapors from this product may injure susceptible plants in the vicinity. Do not store in freezing temperatures. Do not use or store near heat or open flame.

DIRECTIONS FOR USE
EASY TO KILL WEEDS -- Apply when weeds are young and actively growing. 1/4

Use 5 to 10 gal. WKS-65 per 100 gal. of fuel oil or kerosene. Spray a band 12 to 20 inches wide around base of tree to point of run-off. Spray exposed roots of freshly cut stumps to point of run-off to prevent resprouting of undesirable trees. Hard to kill woody plants may require repeat applications if regrowth appears. Use same mixture for stump treatment, and thoroughly wet all exposed areas including roots to prevent regrowth. NOTE: Keep spraying tank agitated.

WKS-65 Controls the Following Weeds:
Cotton Wood, Aspens, Wild Gooseberries, Blackberries, Raspberries and Related Brambles, Oaks, Poison Ivy, Chokecherry, Sassafras, Wild Roses, Sweetgum, Willows, Locust, Perennial Ground Cherry, Horsetail, Currants, Elderberry, Wild Grapes, Plums, Coyote Bush, Mild Vetch, Sumacs, Maples, Elm, Southern Waxmyrtle, Mesquite, Osage Orange, Blackgum, Ground Ivy, Yucca, Prickly Pear, Knotweed.

Herbaceous Broadleaf Weeds Such as:
Cocklebur, Croton weed, Broomweed, Lambsquarter, Curly Indigo, Mexican Weed, Foxglove, Ragweed, Sunflower, Plantains, Chickweed, Dandelions, Oxalis, plus many other species.

CAUTION: Avoid contact with skin, eyes or clothing. Do not take internally. Local conditions may affect the use of WKS-65. Consult state agricultural extension or experiment station weed specialists for specific recommendations on possible local weed problems and for information.

ACCEPTED

JAN 1970