


ABCO ABCO ABCO



USDA REG.
No. 4828-64

MANUFACTURING CHEMISTS

WKC-5

For Outstanding Control of Aquatic Weeds.
A non-selective vegetation killer to control
broadleaf weeds and grasses through contact.

ACTIVE INGREDIENTS:

Diquat dibromide (6,7-Dihydrodipyrido
(1,2-a:2:1'-c) pyrazinediium dibromide)* 2.383%

INERT INGREDIENTS: 97.617%

Total 100.00%

Contains 0.106 pounds cation per gallon

WARNING
KEEP OUT OF REACH OF CHILDREN
(See back panel for additional cautions)

CONTENTS _____ GALS.

4828-64
ACCEPTED
Jul 7 1970
UNDER THE FEDERAL INSECTICIDE
ACT AND RODENT CONTROL
ACTS OF 1947 AND 1958
EPA REG. NO. _____ SUBJECT
ID-ATTACHED COMMENTS

WKC-5

Outstanding for Control of Aquatic Weeds,
also excellent for general weeds in non-crop
areas.

WKC-5 works rapidly and effects are usual-
ly visible within a few days. Do not use on
lawns since this product will kill or top-
kill lawn grasses.

feet apart. If submersed weed growth has
reached the water surface, treat as above
or inject WKC-5 below the surface in strips
no more than 30 feet apart. If algae is also
present, a pre-treatment with Copper Sul-
phate as per recommendation will give
best results with WKC-5.

WARNING
Do not inhale. Do not get on skin. Do
not take internally. May be fatal if
swallowed, inhaled or absorbed through
skin. Prolonged skin contact will cause
severe irritation. Do not get on skin.

NET WEIGHT INGREDIENTS: Total 100.00%
 Contains 0.100 pounds cation per gallon

WARNING
KEEP OUT OF REACH OF CHILDREN
 (See back panel for additional cautions)

CONTENTS _____ GALS.

4828-64
ACCEPTED
 Feb 7 1970
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 THIS PRODUCT IS A POISON BEARING
 THE FOLLOWING NO. _____ SUBJECT
 TO APPLICABLE COMMENTS.

WKC-5

Outstanding for Control of Aquatic Weeds, also excellent for general weeds in non-crop areas.

WKC-5 works rapidly and effects are usually visible within a few days. Do not use on lawns since this product will kill or top-kill lawn grasses.

WEED CONTROL IN NON-CROP OR NON-PLANTED AREAS: WKC-5 will kill annual weeds and topkill perennial weeds such as undesirable grassy and broadleaved weeds around buildings, walking paths, patios, fence lines, and courtyards, park areas, driveways, and similar areas where needed. This product kills by weed contact. The best results are obtained if applied to young weeds. Mature or established weeds may require retreatment for control. Do not spray or allow contact with foliage of crops or ornamental plants.

Use WKC-5 at a rate of 1.85 gals. per 20 gals. of water (0.0925 lbs. per gal. water) for thorough weed contact and weed control.

AQUATIC WEEDS: WKC-5 can be used for both submersed and floating weeds in water, and also marginal weeds along the edges of bodies of water. Use only on weeds in still bodies of water such as ponds, lakes, ditches, laterals and waterways.

SUBMERSED WEEDS	Gals./Surface Area
Bladderwort (<i>Utricularia</i> spp.)	18.5 - 37 gal
Coottail (<i>Ceratophyllum demersum</i>)	37 gal
Elodea (<i>Elodea</i> spp.)	37 gal
Naiad (<i>Najas</i> spp.)	18.5 gal
Pondweeds (<i>Potamogeton</i> spp.)	37 gal
Watermilfoil (<i>Myriophyllum</i> spp.)	18.5 - 37 gal

APPLICATION FOR SUBMERSED WEED CONTROL: Apply at rates recommended above. If weeds have not yet reached the surface, apply WKC-5 from a boat directly from a container into the water while moving slowly over the surface. Pour evenly over area to be treated in strips about 40

feet apart. If submersed weed growth has reached the water surface, treat as above or inject WKC-5 below the surface in strips no more than 20 feet apart. If algae is also present, a pre-treatment with Copper Sulfate as per recommendation will give best results with WKC-5.

MARGINAL WEEDS: For weeds infesting the edges and other portions of ponds, lakes, ditches, laterals, canals and waterways - apply as follows: Cattails (*Typha* spp.) 18.5 gal/100 gal of water.

APPLICATION: For top-kill, apply as a drenching spray for complete coverage and thorough weed contact. Repeat if necessary due to regrowth. For best results apply before flowering.

FLOATING WEEDS	Gals./Surface Area
Pennywort (<i>Hydrocotyle umbellata</i>)	9 1/2 - 13 1/2 gal
Salvinia (<i>Salvinia rotundifolia</i>)	9 1/2 - 13 1/2 gal
Waterhyacinth (<i>Eichhornia crassipes</i>)	9 1/2 - 13 1/2 gal
Waterlettuce (<i>Pistia stratiotes</i>)	9 1/2 - 13 1/2 gal
Duckweed (<i>Lemna</i> spp.)	18.5 gal

APPLICATION FOR FLOATING WEED CONTROL: Apply recommended amounts of WKC-5 above in 8 to 12 gals of water per acre. Make sure you thoroughly cover all plants on water surface and on damp marginal areas. Regrowth of duckweed may occur from untreated areas so that retreatment may be required for season long control.

ALGAE: WKC-5 controls certain filamentous algae. Apply as with submersed weeds. *Spirogyra* spp. and *Pithophora* spp. - 1/2 to 1 1/2 PPM.

Note: In following, PPM refers to cation. Formula For Parts Per Million (PPM)

RECOMMENDATIONS
 Gallons to use = p.p.m. recommended x acreage of water x average depth of water x 25.2.

WARNING

Do not inhale. Do not get on skin. Do not take internally. May be fatal if swallowed, inhaled or absorbed through skin. Prolonged skin contact will cause severe irritation. Do not get on skin, eyes or clothing. Repeated contact with skin may increase danger of absorption. Symptoms of injury may be delayed. In case of skin contact, wash immediately with water; remove clothing and wash skin where necessary. For eyes, wash thoroughly with water and get medical attention. Do not breathe spray mist in order to avoid nasal, throat and respiratory tract irritations. When spraying or when contacting vegetation sprayed previously, wear waterproof footwear and clothing. Do not contaminate feed, food-stuffs or drinking water. Do not apply under conditions where spray can drift to food, forage or other plantings that may be damaged or the crops rendered unfit for sale or consumption. Store above 32°F. Wash and destroy empty container. Do not re-use.

NOTE: Do not use treated water for human or animal consumption, swimming, spraying or irrigation within 10 days after treatment. Since treatment of dense weed areas can result in oxygen loss due to decomposition of dead weeds and thus cause suffocation of fish, treat only part of dense weed areas at a time to prevent this occurrence. Do not apply to muddy water or avoid creating muddy water during application. Do not apply under conditions of high winds and high waves or do not apply to plants covered with mud deposits.

Consult your State Fish and Game Agency before applying this product.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ABCO, INC. P. O. BOX J IRWIN, PA. 15642 U. S. A.
 1209