


ABCO ABCO ABCO

PA EST 4828 PA 1

EPA Reg. No. 4828-64



MANUFACTURING CHEMISTS

WKC-5

For Outstanding Control of Aquatic Weeds. A non-selective
vegetation killer to control broadleaf weeds and grasses
through contact.

ACTIVE INGREDIENTS:

Diquat dibromide (6,7-Dihydrodipyridinyl, 1,2-bis(4-
pyridin-2-yl)ethane) dibromide 2.383%

INERT INGREDIENTS

97.617%

Total

100.000%

*Contains 0.108 pounds cation per gallon

WARNING

KEEP OUT OF REACH OF CHILDREN

(See sig. panel for additional
precautionary statements)

NET CONTENTS _____ GALS

Manufactured by

**ABCO INC. 230 INDUSTRY BLVD.
NORTH HUNTINGDON, PA. 15642**

WKC-5

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

"WARNING"

NOTE: Consult your State Fish and Game Agency before applying this product. Do not inhale. Do not get on skin. May be fatal if swallowed, inhaled or absorbed through skin. Causes skin irritation. Do not get on skin, eyes or clothing. Symptoms of injury may be delayed.

In case of skin contact, wash immediately with water, remove clothing, and wash skin where necessary. For eyes, wash thoroughly with water and get medical attention. If swallowed, drink 1 or 2 glasses of water. Induce vomiting by placing finger in back of throat. Call a physician. Never give anything by mouth to an unconscious person. Do not breathe spray mist in order to avoid nasal, throat and respiratory tract irritations. When spraying or when contacting vegetation sprayed previously, wear waterproof footwear and clothing. Do not contaminate feed, foodstuffs or drinking water.

ENVIRONMENTAL HAZARD

Do not apply directly to water except as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes. Since treatment of dense weed areas can result in oxygen loss due to decomposition of dead weeds and thus cause suffocation of fish, treat only part of dense weed areas at a time to prevent this occurrence.

STORAGE AND DISPOSAL

Storage: Store above 32° F.

Disposal: Triple rinse (or equivalent).

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures. Then either for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved state and local procedures.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For application only to ponds, lake, and slough waters where there is little or no outflow of water and which are

totally under the control of the product user. Do not use the treated water for animal consumption, swimming, spraying, irrigation of domestic purposes for 14 days after treatment. Do not apply to aquatic sites in Florida. Only for sale to, use and storage by Service persons.

USE PRECAUTIONS

Do not apply to muddy water or avoid creating muddy water during application. Do not apply under conditions of high winds and high waves or do not apply to plants covered with mud deposits. Do not apply under conditions where spray can drift to food, forage or other plantings that may be damaged or the crops rendered unfit for sale or consumption. Store above 32° F.

WKC-5 works rapidly and effects are usually visible within a few days. Do not use on lawns since this product will kill or top-kill lawn grasses.

WEED CONTROL IN NONCROP OR NONPLANTED AREAS: WKC-5 will kill annual weeds and top-kill perennials weeds such as undesirable grassy and broadleaved weeds around buildings, walking paths, patios, fence lines, golf courses, park areas, driveways, and similar areas where needed. This product kills by weed contact. The best results are obtained if applied to young weeds. Mature or established weeds may require treatment for control. Do not spray or allow contact with foliage of food crops or ornamental plants.

Use WKC-5 at rate of 1.85 gals. per 20 gals. of water (1.85 gal. per gal. water) for thorough weed contact and weed control.

AQUATIC WEEDS: WKC-5 can be used for both submersed and floating weeds in water, and also marginal weeds along the edges of bodies of water. Use only on weeds in still bodies of waters such as ponds, lakes and ditches.

SUBMERSED WEEDS

	Gal. Surface Acre
Bladderwort (<i>Utricularia</i> spp.)	18.5/32 gal.
Coontail (<i>Ceratophyllum demersum</i>)	37 gal.
Elodea (<i>Elodea</i> spp.)	37 gal.
Najas (<i>Najas</i> spp.)	18.5 gal.
Pondweeds (<i>Potamogeton</i> spp.)	37 gal.
Watermilfoil (<i>Myriophyllum</i> spp.)	18.5/32 gal.

APPLICATION FOR SUBMERSED WEED CONTROL: apply at rates recommended above. If weeds have not yet reached the surface, apply WKC-5 from a boat directly from a container into the water while moving slowly over the surface.

Post evenly over area to be treated in strips about 40 feet apart. If submersed weed growth has reached the water surface, treat as above or inject WKC-5 below the surface in strips no more than 20 feet apart. If algae is also present, a pretreatment with Copper Sulphate as per recommendation will give best results with WKC-5.

MARGINAL WEEDS: For weeds infesting the edges and other portions of ponds, lakes and ditches, apply as follows: Cattails (*Typha* spp.) 18.5 gal./100 gal. of water.

APPLICATION: For top-kill, apply as a drenching spray for complete coverage and thorough weed contact. Repeat if necessary due to regrowth. For best results apply before flowering.

FLOATING WEEDS

	Gal. Surface Acre
Pennyworth (<i>Hydrocotyle umbellata</i>)	9 1/2-13 3/4 gal.
Salvinia (<i>Salvinia Rotundifolia</i>)	9 1/2-13 3/4 gal.
Waterhyacinth (<i>Echhornia crassipes</i>)	9 1/2-13 3/4 gal.
Waterlettuce (<i>Pistia stratiotes</i>)	9 1/2-13 3/4 gal.
Duckweed (<i>Lemna</i> spp.)	18.5 gal.

APPLICATION FOR FLOATING WEED CONTROL: Apply recommended amount of WKC-5 above in 8 to 12 gals. of water per acre. Make sure you thoroughly cover all plants on water surface and on damp marginal areas. Regrowth of duckweed may occur from untreated areas so that retreatment may be required for season long control.

ALGAE: WKC-5 controls certain filamentous algae. Apply as with submersed weeds. *Spirogyra* spp. and *Pithophora* spp. to 1% PPM.

NOTE: In following, PPM refers to cacon. Formula for Parts Per Million (PPM).

RECOMMENDATIONS

Gallons to use = ppm recommended x acreage of water x average depth of water x 2.52.

Labeler makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ABCO, Inc. POST OFFICE BOX J • IRWIN, PENNSYLVANIA 15642