

IMPEDE

Liquid Growth Retardant For Controlling Grass Growth

Slows down grass growth and reduces number of mowings required. Use only on good grass requiring frequent cutting or areas difficult to cut. Use only on turf that is at least 3 years old.

Best results obtained when plants are growing rapidly and soil is moist. Poor results obtained when soil is dry, leaves are wilted or during extended periods of dryness.

Apply when no rain is expected for at least 12 hours thereafter. After product is absorbed, rain has no effect on performance.

DIRECTIONS

For proper control of grass growth, apply only to green, not brown grass. Apply once either in fall, from October 1st to November 15th, or in the spring from April 10th to June 1st. These are ideal application periods. A mowing thereafter may be required to improve appearance. Do not spray newly mowed areas, as grass clippings may prevent absorption by shorter grass blades. Allow one week after applying before you mow.

Spring application - Use 2 $\frac{2}{3}$ gallons of Impede in 100 gallons of water per acre for good results.

Fall application - Use 3 $\frac{1}{3}$ gallons of Impede in 100 gallons of water per acre to reduce growth next season. Apply to green grass before dormant. Do not apply after leaves have fallen, because they interfere with coverage and absorption, thus results will be poor. Fall application controls wild onion, garlic and biennial type weeds like dandelion and plantain.

Do not apply in summer as permanent grasses become partially dormant.

ANNUAL BLUE GRASS CONTROL - Poa annua on Golf fairways can be reduced. Mow area twice in normal sequence, 5 to 8 days apart. Before first Poa annua seed heads appear, and before next required

mowing, spray 2 quarts of Impede in 30 to 40 gallons of water per acre. Results should be seen in 8 to 10 days and treatment should eliminate or reduce reseeding of Poa annua with little effect on grass growth.

OVERSEEDING RYEGRASS IN BERMUDA GRASS - Excess growth of Bermuda Grass on golf greens in the fall is often a problem. Apply 1 $\frac{1}{2}$ gallons of Impede in 100 gallons of water per acre in late September or early October to inhibit growth and enable rye or other grasses to produce a good stand. The Bermuda grass must be in good condition. Overseed no sooner than 48 hours after spraying.

INHIBITING GRASS ALONG EDGES OF HOME LAWNS - Apply the equivalent of 2 $\frac{2}{3}$ gallons of Impede in 200 gallons of water per acre between April 10th and June 1st to a 6 inch band along sidewalks and edges of shrubs and trees. Spray a 6 inch band 800 feet long with a compressed air shoulder sprayer using 8 tablespoonfuls in 2 gallons of water.

CAUTION!


Use only for grass inhibition. Do not pasture livestock, or cut hay from treated areas. Use only along highways, airports, golf course greens, fairways and similar areas. Only use along edges of home lawns. No effect on grass color at proper application rates.

Container disposal - Chop holes in top and bottom and bury 18 inches below surface at safe disposal site. Never reuse for food or water.

Fish and Wildlife - Do not contaminate water by cleaning of equipment or disposal of wastes.

Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instruction.

ABCO INC. • R. D. 3 IRWIN, PA. 15642



IMPEDE

Liquid Growth Retardant For Controlling Grass Growth

Slows down grass growth and reduces number of mowings required. Use only on good grass requiring frequent cutting or areas difficult to cut. Use only on turf that is at least 3 years old.

Best results obtained when plants are growing rapidly and soil is moist. Poor results obtained when soil is dry, leaves are wilted or during extended periods of dryness.

Apply when no rain is expected for at least 12 hours thereafter. After product is absorbed, rain has no effect on performance.

DIRECTIONS

For proper control of grass growth, apply only to green, not brown grass. Apply once either in fall, from October 1st to November 15th, or in the spring from April 10th to June 1st. These are ideal application periods. A mowing thereafter may be required to improve appearance. Do not spray newly mowed areas, as grass clippings may prevent absorption by shorter grass blades. Allow one week after applying before you mow.

Spring application - Use 2 2/3 gallons of Impede in 100 gallons of water per acre for good results.

Fall application - Use 3 1/2 gallons of Impede in 100 gallons of water per acre to reduce growth next season. Apply to green grass before dormant. Do not apply after leaves have fallen, because they interfere with coverage and absorption, thus results will be poor. Fall application controls wild onion, garlic and biennial type weeds like dandelion and plantain.

Do not apply in summer as permanent grasses become partially dormant.

ANNUAL BLUE GRASS CONTROL - Poa annua on Golf fairways can be reduced. Mow area twice in normal sequence, 5 to 8 days apart. Before first Poa annua seed heads appear, and before next required

mowing, spray 2 quarts of Impede in 30 to 40 gallons of water per acre. Results should be seen in 8 to 10 days and treatment should eliminate or reduce reseeding of Poa annua with little effect on grass growth.

OVERSEEDING RYEGRASS IN BERMUDA GRASS - Excess growth of Bermuda Grass on golf greens in the fall is often a problem. Apply 1 1/2 gallons of Impede in 100 gallons of water per acre in late September or early October to inhibit growth and enable rye or other grasses to produce a good stand. The Bermuda grass must be in good condition. Overseed no sooner than 48 hours after spraying.

INHIBITING GRASS ALONG EDGES OF HOME LAWNS - Apply the equivalent of 2 2/3 gallons of Impede in 200 gallons of water per acre between April 10th and June 1st to a 6 inch band along sidewalks and edges of shrubs and trees. Spray a 6 inch band 800 feet long with a compressed air shoulder sprayer using 8 tablespoonfuls in 2 gallons of water.

CAUTION!

Use only for grass inhibition. Do not pasture livestock, or cut hay from treated areas. Use only along highways, airports, golf course greens, fairways and similar areas. Only use along edges of home lawns. No effect on grass color at proper application rates.

Container disposal - Chop holes in top and bottom and bury 18 inches below surface at safe disposal site. Never reuse for food or water.

Fish and Wildlife - Do not contaminate water by cleaning of equipment or disposal of wastes.

Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ABCO INC. • R. D. 3 IRWIN, PA. 15642