

This product is recommended for control of:

**BLACK SPOT and POWDERY MILDEW on ROSES**

**POWDERY MILDEW on CHRYSANTHEMUMS, CALENDULA, DAHLIAS, PHLOX, SWEET PEAS, DELPHINIUMS, LILACS, BEANS, PEAS, PEACHES**

**RUST on CHRYSANTHEMUMS, SNAP DRAGONS (before or after blossoming), CARNATIONS, ASPARAGUS and BEANS**

**COMMON RED SPIDER, PACIFIC MITE on BEANS, COTTON, EVERGREENS, FIGS, PEAS, PLUMS**

**THRIPS, CITRUS RED SPIDER and TWO SPOTTED MITE on CITRUS FRUITS**

Apply early in the morning or cool of the evening when the air is still. Thoroughly cover all stems, leaves and blossoms on both upper and lower surfaces. As a fungicide, frequent applications should be made early in the season. During the summer and fall, applications should be made at 10 day intervals. As an insecticide, apply carefully when the insects threaten and repeat every few days until the pests are under control.

### USE AS A SPRAY

3 or 4 tablespoons to 1 gallon of water. Add slowly with continuous stirring until sulfur is thoroughly mixed. Apply with ordinary sprayer. Keep mixture well stirred at all times.



### USE AS A DUST

Use this WETTABLE DUSTING SULFUR as it comes from this can or apply with a hand duster.

### CHIGGER CONTROL

Control chiggers in the yard by dusting over the entire lawn, flower garden, shrubs and any place the chiggers may be hiding at weekly intervals or a day or two before protection is desired. The contents of this container will cover from 750 to 1,000 square feet.

**CONTAINER DISPOSAL:** Do not reuse empty container. Destroy it by perforating or crushing. Bury in a safe place away from water supplies.



# DUST

US

## CONTROLS:

**ACTIVE INGREDIENT**

Sulfur .....

**INERT INGREDIENTS** .....

93% of the Sulfur

PACKED BY  
STEPHENSON CHEMICAL CO., INC.  
COLLEGE PARK, GEORGIA

STEPHE