

BACTIN

BACTIN Bacteriostat-Sanitizer is based on a combination of quaternary ammonium compounds.

BACTIN is designed for laundry use in the final rinse cycle. However, already laundered fabric may be treated by soaking in a BACTIN solution as recommended under Directions.

BACTIN provides effective residual bacteriostatic and self-sanitizing properties for laundered items, such as athletic equipment, hospital and institutional linens and diapers.

BACTIN is used in hospitals, nursing homes, hotels, motels, restaurants, schools and commercial laundries.

NOTE: For residual bacteriostatic or selfsanitizing activity, conditions of high humidity or wetness are required. BACTIN is concentrated, thus economical in use. Handle with care.

DIRECTIONS FOR USE

Use 2.24 fluid ounce (14 teaspoons) per 100 lbs. of dry fabric for residual bacteriostatic or self-sanitizing activity against Klebsiella Pneumoniae, aberrant, ATCC No. 4352 and Staphylococcus aureus ATCC No. 6538.

Dilute recommended amount of BACTIN in ¼ to ½ gallon of water; then add to the wash wheel in the final rinse. Retreat fabrics after each washing.

Do not reuse empty drums. Return to drum reconditioner or rinse well with soap solution and discard. Do not contaminate water by disposal of waste.

DANGER!

KEEP OUT of REACH of CHILDREN.


Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Huyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ABCO, INC. 230 Industry Boulevard, North Huntingdon, Pa. 15642



BACTIN

LIQUID BACTERIOSTAT - SANITIZER FOR INSTITUTIONAL AND INDUSTRIAL LAUNDRIES

Acuve Ingredients	
Octyl decyl dimethyl ammonium chloride Dioctyl dimethyl ammonium chloride Didecyl dimethyl ammonium chloride Isopropyl alcohol	3.125%
Inert Ingredients	5.000%
Water	82.500%
	100.00%

DANGER!

KEEP OUT of REACH of CHILDREN

(See back panel for first aid statement and other precautions)

NET CONTENTS ____ GALS.

374

CCEPTED 3-14-74 4828-95

THE FEDERAL INSECTICIDE FUNCTION AND ROBENTICIDE ACT FOR CONOMIC POISON REGISTER-ED U. DER NO.

BACTIN