


EPA. EST. No. 4828-PA-1

EPA. REG. No. 4828-102



4828-102

MANUFACTURING CHEMISTS

CLASH

RAPID CONTROL OF FLEAS AND TICKS
ON CATS AND DOGS
Provides Residual Control Also
ODORLESS - NON-STAINING

Contains DRIONE and SEVIN[®] Insecticides

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
Piperonyl Butoxide, Technical	1.00%
Carbaryl (1-Naphthyl N-methylcarbamate)	5.00%
Silica Gel	40.00%
Base Oil	49.90%

INERT INGREDIENTS

49.00%
Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and to 0.2% of related compounds.

CAUTION!

KEEP OUT of REACH of CHILDREN

(See rear panel for additional precautions)

NET CONTENTS _____ POUNDS
1174

CLASH

Flea and Tick Insecticide Powder

Apply liberally to body of animal and work in dust by rubbing against the lay of the hair until it penetrates to the skin.

Repeat once weekly if needed. Contact ticks with powder.

Do not treat kittens or puppies under 4 weeks of age.

For kennels and living quarters, dust bedding and floor at the rate of 1 ounce per 50 square feet of surface. Repeat as needed.

4828-102

CAUTION!

Harmful if swallowed or inhaled. Avoid contamination of feed and foodstuffs or inhalation of dust. Avoid contact with skin. Wash hands thoroughly after use.

Carbaryl (1-Naphthyl N-methylcarbamate)	1.00%
Silica Gel	5.00%
Base Oil	40.00%
INERT INGREDIENTS	49.00%
Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and to 0.2% of related compounds.	

CAUTION!
KEEP OUT of REACH of CHILDREN
 (See rear panel for additional precautions)

NET CONTENTS _____ POUNDS
 1174

CLASH

Flea and Tick Insecticide Powder

Apply liberally to body of animal and work in dust by rubbing against the lay of the hair until it penetrates to the skin.

Repeat once weekly if needed. Contact ticks with powder.

Do not treat kittens or puppies under 4 weeks of age.

For kennels and living quarters, dust bedding and floor at the rate of 1 ounce per 50 square feet of surface. Repeat as needed.

4828-102

CAUTION!

Harmful if swallowed or inhaled. Avoid contamination of feed and foodstuffs or inhalation of dust. Avoid contact with skin. Wash hands thoroughly after use.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on this label.

Do not reuse container. Destroy by perforating or crushing and burying in a safe place.

Drione[®] is a registered trademark of FMC Corporation.

*Sevin is a trademark of Union Carbide Corporation for the active ingredient Carbaryl.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ABCO INC.
230 Industry Boulevard
North Huntingdon, Pa. 15642

STOP! READ THE LABEL

POISON

See Side Panels for Antidote & Precautions



CAN KILL YOU

DANGER

Keep Out Of Reach of Children

ATOMIC

METHYL PARATHION 4E

(Contains 4 lb. Methyl Parathion per Gallon)

NOT FOR HOME USE

ACTIVE INGREDIENTS:

*O, O-dimethyl O-p-nitrophenyl thiophosphate	45.96%
Xylene	48.89%
INERT INGREDIENTS	5.15%
TOTAL	100.00%

*Methyl Parathion

E.P.A. REG. NO. 4977-107

SOUTHEASTERN Chemical CORP.

Fairfax, South Carolina - EPA EST NO. 4977-SC-1

Bishopville, South Carolina - EPA EST. NO. 4977-SC-2

NOTICE OF WARRANTY

Southeastern Chemical Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions or Use and Conditions of Sale, subject to the inherent risks referred to herein. Southeastern Chemical Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY, and no agent of Southeastern Chemical Corporation is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and not include consequential commercial damages such as loss of profit or value, etc.

DIRECTIONS FOR USE

Mix in water with agitation; agitate while spraying. Apply to row and field crops in 3 to 10 gals. (California only: use 20 to 100 gals.) of water per acre by ground spray rig or 1 to 3 gals. (California only: use 5 to 15 gals.) of water by airplane.

COMPATIBILITY - Avoid combinations containing highly alkaline materials such as lime-sulfur solutions or lime. Alkaline compounds will reduce the effectiveness of Parathion.

BEANS: Leafhoppers, flea beetles, plant bugs (including Lygus bugs), aphids, armyworms - Apply 1-2 1/2 pints per acre. Cabbage loopers - Apply 2 pints per acre. Stink bugs, cutworms (including climbing and surface feeding species) - Apply 2-2 1/2 pints per acre. Do not apply within 15 days of harvest. For green and lima beans, do not apply within 21 days of harvest for dosage rates exceeding 1 pint per acre.

CABBAGE: Aphids, armyworms, thrips - Apply 1/2-2 pints per acre. Flea beetles, mites - Apply 1-2 pints per acre. Imported cabbageworms, cabbage loopers, stink bugs - Apply 2-2 1/2 pints per acre. Plant bugs - Apply a minimum dosage of 3 pints of product per acre. Do not apply within 21 days of harvest.

COLLARDS: Aphids, armyworms, flea beetles, leafhopper, mites - Apply 1-2 pints per acre. Imported cabbageworm, cabbage looper, stink bugs, thrips - Apply 2-3 pints per acre. Plant bugs - Apply 3 pints per acre. Do not apply within 21 days of harvest.

CORN: Aphids (including corn leaf aphid), armyworm, fall armyworm, corn rootworm, (including western corn rootworm) (adults), flea beetles, mites, stink bugs - Apply 1/2 pint per acre. Do not apply within 12 days of harvest. Workers entering treated fields within 24 hours after application should wear protective clothing.

COTTON: Thrips, (apply before or during the 4-leaf stage and repeat 1 to 3 times at 7 to 10 day intervals), fleahoppers (including cotton fleahopper) - Apply 1/2 to 1 pint per acre. Grasshopper - Apply 1/2 pint per acre. Cotton leafworm, garden webworm - Apply 1/2-1 pint per acre. Lygus bugs, spider mites (including desert spider mite and two-spotted spider mites) - Apply 1/2-2 pints per acre. Southern green stink bug, false chinch bug - Apply 1 pint per acre. Cabbage looper, cotton leaf-perforator, cutworms, fall armyworm (to 3rd instar), aphids (including cotton aphid) (apply when aphids appear in buds and on leaves of seedlings or when honeydew appears on older plants) - Apply 1-1 pints per acre. Leafrollers - Provide a minimum dosage of 1.2 pints of product per acre. Saltmarsh caterpillar, stink bugs - Apply 2 pints per acre. Armyworm - Apply 2-4 pints per acre. Bollworm - Apply 2-6 pints per acre. Do not handpick or harvest within 7 days of application. Workers entering fields within 24 hours after application should wear protective clothing. Boll Weevil - Apply 1-6 pints per acre. For early season control of over-wintering boll weevils, apply as early as the 2 to 4-leaf stage and repeat 1-3 times at 7 to 10 day intervals. For mid season control, apply at the start of squaring or after 10-25% of the first squares are punctured and repeat at 3 to 8 day intervals as needed or until the top bolls are full grown. Repeat at 3-day intervals during the major fruiting season, and periods of heavy weevil migration.

CUCUMBERS: Aphids, mite, two-spotted spider - Apply 1/2 pint per acre. Do not apply before vining. Do not apply within 15 days of harvest.

GRAPES: Mites, thrips - Apply 1 1/2 pints per acre. Aphids, grape leafroller, grape leafhopper - Apply 1/2-2 pints per acre (per 100 gal. water). Do not apply within 14 days of harvest.

GRASS (HAY, PASTURE): Armyworm (to 3rd instar), black grass bugs (including Irbisia), false chinch bug, grasshopper, leafhopper - Apply 1 1/2 pints per acre. Do not apply within 15 days of harvest.

LETTUCE: Aphids, armyworms, flea beetles, imported cabbageworm, leafhoppers, mites - Apply 1-2 pints per acre. Cabbage looper, plant bugs - Apply 2 pints per acre. Do not apply within 21 days of harvest.

MUSTARD GREENS: Aphids, armyworm - Apply 1/2-2 pints per acre. Flea Beetles, leafhoppers, mites - Apply 1-2 pints per acre. Imported cabbageworm, cabbage looper, stink bugs - Apply 2-3 pints per acre. Plant bugs - Apply 3 pints per acre. Do not apply within 10 days of harvest through 1 pint, or within 21 days of harvest from above 1 pint through 3 pints.

OATS: Brown wheat mite - Apply 1/4 - 1 pint per acre. Aphids (including greenbug), black grass bugs (including Irbisia) - 1/4-1 1/2 pints per acre. Barley thrips - 1/2 - 1 pints per acre. Climbing cutworms - Provide a minimum dosage of 1/4 pint of product per acre. Leafhoppers - 1/2-1 pint per acre. Armyworm (to 3rd instar), wintergrain mite, stink bugs - 1/2-1 1/2 pints per acre. Say's stink bugs, chinch bugs, false chinch bug - 1 1/2 pints per acre. Do not apply within 15 days of harvest from 1/2 pint through 1 1/2 pints.

PASTURE: (Irrigated) (Refer also to grass): Mosquitoes (larvae) - Apply 1/5 pint per acre. Do not apply within 7 days of harvest. Broadcast application.

PEACHES: Aphids, (including green peach aphid and mealy plum aphid), European red mite - Apply 1/4 - 1 pint per 100 gals. water. Do not apply within 14 days of harvest. Apply in early cover treatment or apply twice within 5 day intervals as needed. Oriental fruit moth, peachtree borer, plum curculio - Apply 1/2 - 2 pints per acre (per 100 gals. water). For peachtree borer apply 2 to 3 sprays to trunks from ground to scaffold limbs. Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

PEAS: Aphids, armyworm (to 3rd instar), flea beetles, leafhoppers, mites, plant bugs (including Lygus bugs) - Apply 1-2 pints per acre. Stink bugs, cowpea curculio, cutworm (including climbing and surface feeding species), cabbage looper - Apply 2 pints per acre. Do not apply within 10 days of harvest through 1 pint. Do not apply within 15 days of harvest from 1 pint through 2 pints.

PEPPERS: Aphids, armyworms, flea beetles, leafhoppers, mites - Apply 1-2 pints per acre. Cutworm (including climbing and surface feeding species), cabbage looper, plant bugs - Apply 2 pints per acre. Do not apply within 15 days of harvest.

POTATOES: Aphids, flea beetles - Apply 1/2-2 pints per acre. Armyworm, leafhoppers, mites - Apply 1-2 pints per acre. False chinch bugs, cabbage looper, plant bugs (including Lygus bugs), stink bugs - Apply 3 pints per acre. Do not apply within 5 days of harvest.

SOYBEANS: Climbing cutworms - Apply 1/2 pint per acre. Three cornered alfalfa hopper, webworm (including garden webworm) - apply 1/2 - 1 pint per acre. Thrips, velvetbean caterpillar - Apply 1/4 - 1 pint per acre. Stink bugs - Apply a minimum dosage of .7 pints of product per acre. Mexican bean beetle, spider mites, aphids, blister beetle - Apply 1 pint per acre. Green cloverworm, cabbage looper - Apply 1 - 2 pints per acre. Armyworm, fall armyworm (to 3rd instar), bean leaf beetle, corn earworm, grasshoppers - Apply 2 pints per acre. Do not apply within 20 days of harvest. Do not apply more than twice per growing season.

STRAWBERRIES: Aphids - Apply 1 - 1 1/2 pints per acre. Mites - Apply 1 1/2 pints per acre. Do not apply within 14 days of harvest.

SWEET POTATOES: Aphids, armyworm (to 3rd instar), cutworms (including climbing and surface feeding species), flea beetles, leafhoppers, cabbage looper, spider mites - Apply 1 1/2 pints per acre. Do not apply within 5 days of harvest.

TOBACCO: Green June Beetles - Apply a minimum dosage of 3/8 pint of product per 100 gal. water to an area of 100 square yards. Apply as a drench to plant bed with a sprinkling can. Do not apply within 5 days of transplanting. Aphids (including green peach aphid), cutworms (including climbing and surface feeding species) - Apply 1/2 - 1 pint per acre. Do not apply within 15 days of harvest. Do not apply within 5 days of priming. Avoid contact with plant juices when priming or cutting tobacco. When necessary to enter treated fields within 24 hours after application, protective clothing should be worn.

TOMATOES: Tomato russet mite - Apply 1/4-1 pint acre. Aphids, armyworm - Apply 1/2-2 pints per acre. Leafhoppers, mites - Apply 1-2 pints per acre. Potato psylla - Apply 1-3 pints per acre. Plant bugs - Apply 2 pints per acre. Cabbage looper - Apply 2-3 pints per acre. Do not apply within 15 days of harvest from 1 pint through 3 pints.

TURNIPS: Aphids (including cabbage aphid), armyworms (to 3rd instar) - Apply 1/2-1 1/2 pints per acre. Flea beetles, leafhoppers, leafminer, mites - Apply 1-1 1/2 pints per acre. Imported cabbage worm, cabbage looper, plant bugs - Apply 1 1/2 pints per acre. Do not apply within 7 days of harvest through 1/2 pint. (roots) Do not apply within 15 days of harvest from 1/2 to 1 1/2 pints (roots). Do not apply within 21 days of harvest if tops are to be used as food or feed.

WHEAT: Brown wheat mite - Apply 1/4-1 pint per acre. Barley thrips - Apply 1/2-2/3 pints per acre. Leafhoppers - Apply 1/2-1 pint per acre. Climbing cutworms - Apply 3/4-1 pint per acre. Aphids (including greenbug), black grass bugs (including Irbisia) - Apply 1/4-1 1/2 pints per acre. Winter grain mite, stink bug, armyworm (to 3rd instar) - Apply 1/2-1 1/2 pints per acre. Say's stink bugs, chinch bug, false chinch bug - Apply 1 1/2 pints per acre. Do not apply within 15 days of harvest from 1/2 pint through 1 1/2 pints.

ORNAMENTALS: (Woody Shrubs, Trees and Vines), Christmas tree plantation (Pines) - European pine shoot moth, Nantucket pine tip moth - Apply 2 pints per acre. Apply in 5 to 50 gallons of water.

FOREST, NONAGRICULTURAL LAND, WASTELANDS: Mosquitoes (larvae) - Apply 1/5 pint per acre. Do not apply within 7 days of harvest through 1/5 pint. For use by mosquito abatement districts and official government agencies only. Repeat as needed. Toxic to fish and wildlife. Do not use where shrimp and crabs are an important resource. Do not use in any estuaries, or coastal or tidal areas along the southeast Atlantic or Gulf coasts of the United States.

PINE FORESTS: European pine shoot moth, Nantucket pine tip moth - Apply 2 pints per acre. Apply in 5 to 50 gallons of water.

SOUTHEASTERN CHEMICAL CORP.

P. O. Box 1026

Orangeburg, S. C. 29115

NET CONTENTS

U.S. GALLONS

ROUSSEL

METALDEHYDE

ACTIVE INGREDIENT

metalddehyde minimum

INERT INGREDIENTS up to

TOTAL

100

100

100

DIRECTIONS:

FOR MANUFACTURING AND REFORMULATING USES ONLY

WARNING

KEEP OUT OF REACH OF CHILDREN

May be harmful or fatal if swallowed. Avoid breathing dust and vapor. Use with adequate ventilation. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. In case of contact, flush with plenty of water. For eyes, get medical attention. Avoid contamination of food and feedstuffs with this material.

This material is toxic to fish and other aquatic life. Do not discharge into any body of water without proper consultation with appropriate authorities.

Do not use in empty containers. Do not use in containers that have previously contained a different material. Do not use in containers that have previously contained a different material.

Keep away from children and animals. Keep away from food and feedstuffs.

For more information:

Roussel Corporation
155 East 44 Street
New York, N.Y. 10017 (U.S.A.)



NET WEIGHT

Kilos

Pounds

EPA Reg. No. 5086-9

