

[48231]

**CAUTION**

KEEP OUT OF REACH OF CHILDREN  
Harmful if swallowed.

CALL PHYSICIAN IMMEDIATELY. See an  
antidote. Avoid spraying directly on foodstuffs.  
Do not apply directly on very young animals.  
Do not spray into or near fire or open flame.  
Do not use in poultry houses.

Do not apply as a space spray in a food pro-  
cessing plant while plant is in operation. When  
applied as a surface or contact spray in a food  
processing plant remove or protect food dur-  
ing treatment. After treatment wash with  
potable water all equipment, benching, shelv-  
ing etc.

Do not reuse empty drum. Return to drum  
reconditioner or destroy by perforating or  
crushing and burying in a safe place.

This product is toxic to fish. Keep out of  
lakes, streams, or ponds. Apply this product  
only as specified on this label.

**NOTICE**

Buyer assumes all risks of use, storage, or  
handling of this material not in strict accord-  
ance with directions given herewith.

**ANTIDOTE**

**EXTERNAL:** Wash with cool soapy water.  
**INTERNAL:** Administer soap solution to in-  
duce vomiting. Administer vegetable oil to  
reduce gastrointestinal absorption.

**EYES:** Flush with fresh water for at least 15  
minutes.

Call physician

**COASTAL  
FORMULA I**

**Bug Killer**

**DIRECTIONS  
INDOOR SPRAY**

FOR CONTROL OF FLIES, MOSQUITOES,  
GNATS AND ADULT MOTHS in homes, food  
establishments, farm buildings and warehouses;  
Close all windows and stop all ventilators.  
Direct fine mist or fog upward toward the  
center of the room and around the sides until  
the area is filled with fog or mist particles.  
Keep area closed for 10 to 15 minutes. Clean  
up and destroy fallen insects.

Pilot lights should be extinguished. Cover or  
remove aquariums before treatment. Allow  
only fine mist or "dry" fog to contact fabrics  
or surfaces that might be affected by the oil  
base. Do not remain in treated areas and ven-  
ilate the area after treatment is completed.

FOR CRAWLING INSECTS: ROACHES,  
SILVERFISH, EXPOSED WEAVILS AND  
FLOUR AND GRAIN BEETLES: Set equip-  
ment to deliver a coarse spray. Direct the spray  
to hit as many insects as possible and to cover  
tracing areas to the point of run off and to  
penetrate hiding places. If ANT control is also  
desired, be sure that doors and window cas-  
ings are thoroughly sprayed and that all trail  
ways and nests near the building are treated.  
Check the building's foundations for cracks,  
spraying these cracks and patching them is  
possible.


FOR WASPS AND SPIDERS: Spray insects  
directly and apply heavy spray to nests and  
webs. The most effective wasp control is ob-  
tained by spraying after dark when all insects  
have returned to the nest.

FOR CONTROL OF CARPET BEETLES:  
Clean all rugs, carpets and fabrics to be treated.  
Spray both surfaces of carpets and floor and  
beneath evenly with a fine mist spray. Partic-  
ular attention should be paid to cracks and  
crevices around the baseboards.

ANIMAL SPRAY FOR HORSES AND OTHER  
AND BEEF CATTLE: For protection against  
Stableflies, Hornflies, and Houseflies on the  
animals, spray the animal against the grain of  
the hair, directing a fine mist around the base  
of the horns, on the legs and along the neck  
and belly and over the back. Do not spray  
addres. Apply just enough to wet the ends of  
the hair but not enough to soak the hair. Use  
one or two spray, apply a maximum of 2 fluid  
ounces of spray per animal. Apply once daily  
or as necessary.

NET CONTENTS

DIRECTIONS  
USE OF SPRAY



TO REDUCE THE ANNOYANCE OF FLIES, MOSQUITOES,  
GNATS AND ADULT MOTHS in homes, food estab-  
lishments, farm buildings and warehouses, spray  
indoor areas such as living rooms, dining rooms,  
kitchens, bedrooms, bathrooms, and play areas.  
Close all windows and stop all ventilators. Direct  
fine mist or fog upward toward the center of the  
room and around the sides until the area is filled  
with fog or mist particles. Keep area closed for  
10 to 15 minutes. Clean up and destroy fallen  
insects.

Do not use in poultry houses. Do not spray  
directly on foodstuffs. Do not spray into or near  
fire or open flame. Do not use in poultry houses.  
Do not spray directly on very young animals.  
Do not spray into or near fire or open flame.  
Do not use in poultry houses.

