CHEMICAL PROPERTIES

EASILY SOLUBLE IN ALL POOL WATER LONG LASTING CHLORINE RESIDUAL FORTIFIED WITH WATER SOFTENERS

ACCEPTED 4829-5/ NOV 18 1975

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO.

DANGER:

KEEP OUT OF REACH OF CHIEDREN. Corrosive, causes eye damage. May be fatal if swallowed. Do not get in eyes, on skin or on clothing. Irritating to nose and throat. Avoid breathing dust. May cause burns to broken skin. Wash hands after handling.

FIRST AID:

If swallowed, feed gruel, cooked cereal or bread soaked in milk, followed by olive oil or cooking oil. In case of personal contact, remove excess material and flush skin or eyes with cold water for at least 15 minutes. For eyes get medical attention. Remove and wash contaminated clothing before reuse.

DANGER:

STRONG OXIDIZING AGENT. Mix only with cold water. Use clean, dry utensils. Contamination by moisture, organic matter or other chemicals may liberate hazardous gases. Store in cool, dry, well ventilated area away from heat or open flame. Decomposes at 350°-475°F with liberation of harmful gases. In case of decomposition, if possible, isolate container in open air. Flood with large amounts of water. Keep container tightly closed when not in use. Do not reuse empty container. Rinse empty container thoroughly with water before discarding.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public waters. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

MADE & PRINTED IN U.S.A. E.P.A. Reg. No. 4829-51

MLKJI E.P.A. Est. No. 4829-NJ-1



DIRECTIONS FOR USE

Iso Clor Tablets should be added to pools free of visible algae. Add tablets through a floating dispenser. For initial treatment, use 12 to 24 tablets per 1,000 gallons of pool water. For subsequent treatments, add 4 tablets of Iso Clor per 1,500 gallons of water daily or as needed as determined by test kit to maintain a 1.0 to 1.5 ppm. (parts per million) chlorine residual. Adding 1 tablet will increase the available chlorine by approximately 0.3 ppm. High temperatures, organic contamination, increased bather use, rain and other factors may necessitate more frequent treatments. Using a chlorine and pH test kit, test water frequently to assure proper concertration of chlorine and pH. Maintain pH range of 7.2 to 7.6 at all times. For use in wader and splasher pools, recommended desage is 3 Iso Clor tablets per 750 gallons every day. Maintain a chlorine residual of 10 to 1.5 ppm. as determined by a test kit. 6 tablets weigh approximately one ounce.

D/ KE See

Iso Glor

Chornating tablets





algaericide algericide algaericide algaeri

DANGER:

ded to

dd tab

penser.

to 24

f pool

tments,

1,500

needed

) main-

rts per

Adding

ailable

3 ppm.

ontam-

e, rain

essitate

Ising a

: water

COnceli

aintain

Hirnes.

r pools,

o Clor

'y day.

01 1 1

/ a test

mately

KEEP OUT OF REACH OF CHILDREN
See Antidote and Precautions on Back Panel

ISO CLOR CHLORINE TABLETS DAILY NORMAL MAINTENANCE DOSAGE CHART

POOL SIZE	GALLON CAPACITY	()
8'×2'0''	650	2
10'x30''	14.75	.1
12′x36′′	2550	7
15′x48′′	5300	11;
18′x48′′	7600	21
241×4811	13500	36

DO NOT
MIX WITH ANY
OTHER CHLORINE

CHEMICAL PROPERTIES

EASILY SOLUBLE IN ALL POOL WATER
LONG-LASTING CHLORINE RESIDUAL UNDER THE FEDERAL INSECTICIDE
FORTIFIED WITH WATER SOFTENERS FOR ECONOMIC POISON REGISTER-

KEEP OUT OF REACH OF CHILDREN. Corrosive, causes eye damage. May be fatal if swallowed. Do not get in eyes, on skin or on clothing. Irritating to nose and throat. Avoid breathing dust. May cause burns to broken skin. Wash hands after handling.

FIRST AID: ...

If swallowed, feed gruel, cooked cereal or bread soaked in milk, followed by olive oil or cooking oil. In case of personal contact, remove excess material and flush skin or eyes with cold water for at least 15 minutes. For eyes get medical attention, Remove and wash contaminated clothing before reuse.

DANGER:

STRONG OXIDIZING AGENT. Mix only with cold water. Use clean, dry utensils. Contamination by moisture, organic matter or other chemicals may liberate hazardous gases. Store in cool, dry, well ventilated area away from heat or open flame. Decomposes at 3500-4750F with liberation of harmful gases. In case of decomposition, if possible, isolate container in open air. Flood with large amounts of water. Keep container tightly closed when not in use. Do not reuse empty container. Rinse empty container thoroughly with water before discarding.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public waters. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the

MADE & PRINTED IN U.S.A. E.P.A. Reg. No. 4829-51

E.P.A. Est. No. 4829-NJ-1



DIRECTIONS FOR USE

Iso Clor Tablets should be added to pools free of visible algae. Add tablets through a floating dispenser. For initial treatment, use 12 to 24 tablets per 1,000 gallons of pool water. For subsequent treatments. add 4 tablets of Iso Clor per 1,500 gallons of water daily or as needed as determined by test kit to maintain a 1.0 to 1.5 ppm. (parts per million) chlorine residual. Adding 1 tablet will increase the available chlorine by approximately 0.3 ppm. High temperatures, organic contamination, increased bather use, rain : and other factors may necessitate more frequent treatments. Using a chlorine and pH test kit, test water frequently to assure proper concentration of chlorine and pH. Maintain* pH range of 7.2 to 7.6 at all times. For use in wader and splasher pools, recommended desage is 3 Iso Clor tablets per 750 gallons every day: Maintain a chlorine residual of 1.0. to 1.5 ppm. as determined by a test kit. 6 tablets weigh approximately one ounce.

DAN KEEP O See Anti

..... 9

....

• 6....

Iso Clor.

table

NET WT.: 7% LBS. AVDP. Minimum 744 Tablets COASTA





KEEP OUT OF REACH OF CHILDREN See Antidote and Precautions on Back Panel ACTIVE INGREDIENTS: Sodium Dichloro-s-Triazinetrione

Contains 35% Available Chloring

ISO CLOR CHLORINE TABLETS DAILY NORMAL MAINTENANCE DOSAGE CHART

POOL SIZE	GALLON CAPACITY	TABLETS
8′×20′′	650	2
10'x30"	1475	4
12'x36"	2550	7
15'×48''	5300	15
18'x48"	7600	21
24'×48"	13500	36

DO NOT MIX WITH ANY OTHER CHLORINE