

METEOR BRAND ZINC-COPPER 315

MIXING INSTRUCTIONS

Keep agitator running at all times.

MECHANICALLY AGITATED TANK:

Fill tank half full with water, sprinkle product into water when tank is 75% full, add hydrated lime and fill tank.

BY-PASS AGITATED TANK:

Prepare stock solutions of product and hydrated lime. Fill tank 65% full with water, add product then hydrated lime, continue agitation and fill tank.

Product is corrosive to steel.

DIRECTIONS FOR USE

CITRUS – Central California

To control Brown Rot and Septoria mix 5 pounds of Zinc-Copper 315 with 6 pounds of hydrated lime in 100 gallons of water. Use as a full coverage spray. Apply 500 to 700 gallons of finished spray per acre to mature trees in Fall prior to wet season.

CITRUS – Southern California

To control Brown Rot, mix 3 pounds of Zinc-Copper 315 with 4 pounds of hydrated lime in 100 gallons of water. Use as a skirt spray on soil, trunk and lower 3 to 4 feet of foilage. Apply in Fall just prior to wet season. In Coastal areas or other areas where no copper injury has been found use a mixture of 5 pounds of Zinc-Copper 315 with 6 pounds of hydrated lime in 100 gallons of water as a full coverage spray.

FUNGICIDE

ACTIVE INGREDIENT BY WT.

Copper Sulphate Pentahydrate 37.7%

INERT INGREDIENTS 62.3%


Inert contains 22.0% zinc expressed as elemental and 9.5% copper expressed as elemental. Zinc derived from zinc sulfate, monohydrate. Zinc is for correction of zinc deficiency.

USAGE CAUTIONS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CAUTION

KEEP OUT OF REACH OF CHILDREN



HARMFUL IF SWALLOWED


AVOID BREATHING DUST

AVOID CONTAMINATION OF
FEED AND FOODSTUFFS

Use of this material in a manner other than suggested may produce injury to plants and undesirable results.

Dispose of wastes by burying away from water supply in non-crop sites. Burn empty bags or bury with wastes. Avoid inhaling smoke from burning bags. Avoid drifting of product onto neighboring crop or non-crop areas otherwise injury to animal plant or crops may occur. In California dispose of waste and empty containers in accordance with State and local regulations.

State. Reg. No. 4833-50005-AA



CONDITIONS OF SALE:

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller is authorized to make any other warranty, guarantee or direction concerning this product.
2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of the base price.

Manufactured in U.S.A. by
CHEMICAL & PIGMENT CO.

PITTSBURG, CALIFORNIA 94565

NET WEIGHT _____ LBS.

MIXING INSTRUCTIONS

Keep agitator running at all times.

MECHANICALLY AGITATED TANK:

Fill tank half full with water, sprinkle product into water when tank is 75% full, add hydrated lime and fill tank.

BY PASS AGITATED TANK:

Prepare stock solutions of product and hydrated lime. Fill tank 65% full with water, add product then hydrated lime, continue agitation and fill tank.

Product is corrosive to steel.

DIRECTIONS FOR USE

CITRUS - Central California

To control Brown Rot and Septoria mix 5 pounds of Zinc-Copper 315 with 6 pounds of hydrated lime in 100 gallons of water. Use as a full coverage spray. Apply 500 to 700 gallons of finished spray per acre to mature trees in Fall prior to wet season.

CITRUS - Southern California

To control Brown Rot, mix 3 pounds of Zinc-Copper 315 with 4 pounds of hydrated lime in 100 gallons of water. Use as a skirt spray on soil, trunk and lower 3 to 4 feet of foliage. Apply in Fall just prior to wet season. In Coastal areas or other areas where no copper injury has been found use a mixture of 5 pounds of Zinc-Copper 315 with 6 pounds of hydrated lime in 100 gallons of water as a full coverage spray.


FUNGICIDE

ACTIVE INGREDIENT	BY WT.
Copper Sulphate Pentahydrate	37.7%
INERT INGREDIENTS	62.3%

Contains 37.7% active ingredient. Contains 62.3% inert ingredients. Contains 37.7% active ingredient. Contains 62.3% inert ingredients.

USAGE CAUTIONS


This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.



CONDITIONS OF SALE:

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one other than an agent of Seller is authorized to make any other warranty, guarantee or direction concerning this product.
2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of the base price.

CAUTION



HARMFUL IF SWALLOWED
AVOID BREATHING DUST

AVOID CONTAMINATION OF
FEED AND FOODSTUFFS

Use of this material in a manner other than suggested may produce injury to plants and undesirable results.

Dispose of wastes by burying away from water supply in non-crop sites. Burn empty bags or bury with wastes. Avoid inhaling smoke from burning bags. Avoid drifting of product onto neighboring crop or non-crop areas otherwise injury to animal plant or crops may occur. In California dispose of waste and empty containers in accordance with State and local regulations.

State Reg No 483350005 AA

Manufactured in U.S.A. by

HERMICA B. COMPANY