

Net Content:

U.S. D.A. Reg. No. 4841-38

MICRO BLEND

METHYL

PARATHION 4

Emulsifiable Concentrate

Contains Four Pounds Methyl Parathion Per Gallon

DANGER: KEEP OUT OF REACH OF CHILDREN

See Side Panel for additional Warning Statements and Antidote

ACTIVE INGREDIENTS:

O,O-Dimethyl O-p-Nitrophenyl Thiophosphate*	44.00%
Aromatic Petroleum Distillates	50.00%
INERT INGREDIENTS:	6.00%
TOTAL	100.00%

*O, O-Dimethyl O-p-Nitrophenyl Thiophosphate is otherwise known as Methyl Parathion.

ACCEPTED
9/17/77
4841-38
UNLAWFUL TO REPRODUCE OR TRANSMIT IN ANY MANNER WITHOUT THE WRITTEN PERMISSION OF THE U.S. DEPARTMENT OF AGRICULTURE
ATTACHED COMMENTS
ACT REGISTER SUBJECT



POISON



NON-WARRANTY

SELLER makes no warranty of any kind, expressed or implied, concerning the use of this material other than that indicated on label. Buyer assumes all risk of use or handling of this material when such use and/or handling is contrary to label instructions.

MICRO CHEMICAL COMPANY, INC.
WINNSBORO, LOUISIANA

DIRECTIONS

GRAINS: (Wheat, Oats, and Barley) For control of Greenbug (Aphids) use 1 to 1 pint per acre in sufficient water to give good coverage. Repeat as necessary. Do not apply after heads begin to form.

COTTON: Red Spider Mites, Boll Weevils, Leafworms, Aphids, use 1 pint per acre in sufficient water to cover. Repeat at 3 day intervals. Do not apply within 15 days of handpicking.

MICRO BLEND METHYL PARATHION 4 may be applied by either ground equipment or aircraft. For ground equipment, dilute in sufficient water to give adequate coverage (10 to 20 gallons per acre) depending on amount of vegetation present. For aerial applications, dilute in 3 to 10 gallons of water per acre.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Do not feed the treated plants or trash to dairy animals or animals being fattened for slaughter.

DANGER: POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. DO NOT GET IN EYES OR ON SKIN OR ON CLOTHING. WORKERS ENTERING TREATED FIELDS WITHIN 24 HOURS AFTER METHYL PARATHION APPLICATION SHOULD WEAR PROTECTIVE CLOTHING. WEAR A MASK OR RESPIRATOR OF A TYPE APPROVED BY THE U.S. BUREAU OF MINES FOR PARATHION PROTECTION.

THIS PRODUCT NOT FOR USE OR STORAGE IN OR AROUND THE HOME. KEEP OUT OF REACH OF CHILDREN. KEEP ALL UNPROTECTED PERSONS OUT OF AREA OF OPERATION OR VICINITY WHERE THERE MAY BE DANGER OF DRIFT. VACATED AREAS SHOULD NOT BE RE-ENTERED UNTIL DRIFTING INSECTICIDES AND VOLATILE RESIDUES HAVE DISSIPATED. DO NOT CONTAMINATE FEED OR FOODSTUFFS. AVOID USE DURING FLOWERING AND POLLINATION PERIODS AS THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT OR RESIDUES ON CROPS. PROTECTIVE INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE.

DECONTAMINATE CONTAINER BY WASHING THOROUGHLY WITH LYE SOLUTION, THEN DESTROY TO PREVENT REUSE.

This product is toxic to wildlife. Birds and other wildlife in treated areas may be harmed. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICIANS NOTE AND ANTIDOTE

Methyl Parathion exerts muscarine-like effects—i.e., stimulation of the para-sympathetic nervous system. Repeated use of a para-sympathetic depressant, such as meperidine, to treat a patient's tolerance, is indicated to relieve symptoms. 2-PAM is the antidote for and may be administered by combination with atropine. The antidote is not effective if the patient is comatose. For patients suspected of Parathion



POISON



Reg. No. 4841-38

BLEND

N 4

ate

er Gallon

CHILDREN

nts and Antidote

DIRECTIONS

GRAINS: (Wheat, Oats, and Barley) For control of Greenbug (Aphids) use $\frac{1}{2}$ to 1 pint per acre in sufficient water to give good coverage. Repeat as necessary. Do not apply after heads begin to form.

COTTON: Red Spider Mites, Boll Weevils, Leafworms, Aphids, use $\frac{1}{2}$ pint per acre in sufficient water to cover. Repeat at 3 day intervals. Do not apply within 15 days of handpicking.

MICRO BLEND METHYL PARATHION 4 may be applied by either ground equipment or aircraft. For ground equipment, dilute in sufficient water to give adequate coverage (10 to 20 gallons per acre) depending on amount of vegetation present. For aerial applications, dilute in 3 to 10 gallons of water per acre.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Do not feed the treated plants or trash to dairy animals or animals being fattened for slaughter.

DANGER: POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. DO NOT GET IN EYES OR ON SKIN OR ON CLOTHING. WORKERS ENTERING TREATED FIELDS WITHIN 24 HOURS AFTER METHYL PARATHION APPLICATION SHOULD WEAR PROTECTIVE CLOTHING. WEAR A MASK OR RESPIRATOR OF A TYPE APPROVED BY THE U.S. BUREAU OF MINES FOR PARATHION PROTECTION.

THIS PRODUCT NOT FOR USE OR STORAGE IN OR AROUND THE HOME. KEEP OUT OF REACH OF CHILDREN. KEEP ALL UNPROTECTED PERSONS OUT OF AREA OF OPERATION OR VICINITY WHERE THERE MAY BE DANGER OF DRIFT. VACATED AREAS SHOULD BE RE-ENTERED UNTIL DRIFTING INSECTICIDES AND VOLATILE FUMES ARE COMPLETELY ELIMINATED. CONTAMINATE FEED

4841-38
44.00%
50.00%

