

**CAUTION**  
**KEEP OUT of REACH of CHILDREN**  
(See back panel for additional cautions)

NET CONTENTS      GALLONS

4928-79  
**ACCEPTED**  
APR 7 1972  
UNDER THE FEDERAL INSECTICIDE  
ACT AND FEDERAL FUNGICIDE ACT  
FOR CONTROL OF PESTS IN RESIST  
ED PLANTS AND SUBJECT  
TO ALL APPLICABLE REGULATIONS

## C. G. - 30

A Selective Killer Of Crabgrass when used at our recommended application rates in the presence of Kentucky, Merion Blue, Bermuda and Zoysia grasses.

C. G. - 30 is also selective in the presence of sensitive grasses, such as Fescues and Bentgrasses under certain conditions, such as on cool days and when turf is moist.

Discoloration can occur in Fescue and Bentgrass areas, so therefore a lighter dosage is recommended for satisfactory results.

### APPLICATION DIRECTIONS

**YOUNG CRABGRASS** - Mix 1 pint of C. G.-30 with 10 to 20 gallons of water and apply to 4000 sq. ft. of area. Apply 2 to 3 applications at 7 day intervals.

**BRANCHED CRABGRASS** - Mix 1 pint of C. G. - 30 with 10 to 20 gallons of water and apply to 3000 sq. ft. Apply 2 to 3 applications at 7 day intervals.

**TEMPERATURE** - If over 85 F. when applying, make sure of soil moisture, and use a light dosage of 1 pint in 10 to 20 gallons of water per 5000 sq. ft.

Do not use on St. Augustine Grass, Carpet Grass, Centipede Grass or Dichondra.

### CARE TO BE EXERCISED

1. Do not apply to dry turf during hot dry spells.

2. Restore moisture to the soil by watering turf well for a period of time prior to spraying.
3. Do not apply to newly seeded turf until it has been mowed 3 times.
1. On predominantly Bent or Fescue, use light application rate of 1 pint in 10 to 20 gallons of water per 4000 sq. ft.

### CAUTION

Avoid contact with skin or prolonged breathing of spray mist.

Avoid contamination of feed or food-stuffs. Keep away from children and domestic animals. Keep livestock off treated areas.

Wash hands after using. Do not contaminate water used for domestic purposes. Harmful if swallowed.

Do not reuse empty drum.

Return to drum reconitioner or destroy by perforating or crushing and burying in a safe place.

### ANTIDOTE

If taken internally, induce vomiting and call a doctor at once.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

**ABCO INC.      R. D. 3      IRWIN, PA. 15642 U.S.A.**

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call a physician at once.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

NOTICE: Using of this material in a manner, or at any time other than in accordance with the directions may produce plant injury, excessive residues, or other undesirable results. Micro Chemical Co. makes no express or implied warranties other than specified on the label.

GENERAL INFORMATION: MICRO MONO Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. Its activity is enhanced by addition of a suitable surfactant to the spray solution. It is most effective on young, actively growing weeds at air temperatures above 70° F. Its phytotoxic properties are quickly inactivated on contact with soil.

MIXING INSTRUCTIONS: MICRO MONO Herbicide must be thoroughly dissolved. Fill the spray equipment reservoir about half full with water and add the required amount of MICRO MONO Herbicide and surfactant with agitation. Finish filling the reservoir with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store solution in tank for a prolonged period.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place.

#### DIRECTION FOR USE

COTTON: MICRO MONO Herbicide is useful as a directed application in cotton for the post-emergent control of weeds and grasses such as Johnsongrass, Nutsedge, or Dallisgrass, Cocklebur, Barnyardgrass, Ragweed, Sandbur, and Puncture Vine. It should be applied at the rate of 2 1/2 pints per acre.

Add 6 pints of MICRO MONO Herbicide and 1 to 2 quarts of a suitable surfactant to 100 gallons of water and mix thoroughly. Apply spray solution at 40 gallons per acre. Weed control is most effective at temperatures above

Net Contents 5 Gallons USDA Reg. No. 4841-51

# MICRO MONO

## MSMA Liquid Herbicide

### FOR SELECTIVE POST-EMERGENT WEED CONTROL

#### ACTIVE INGREDIENTS:

Monosodium Acid Methanearsonate\* ----- 51.3%

INERT INGREDIENTS: ----- 48.7%


Total Arsenic (as elemental) all in water soluble form 23.7%

\*Product contains 6.67 lbs. MSMA per gallon or equivalent of 12 lbs. Disodium Methanearsonate Hexahydrate per gallon.

### CAUTION:

## KEEP OUT OF REACH OF CHILDREN!

See Side Panel for additional cautions



Winnsboro, Louisiana 71295

MICRO MONO Herbicide Distributed by Micro Chemical Co., Inc. Winnsboro, La. 71295.

gallon of spray solution per inch of band width per acre. Direct the spray solution to the base of the cotton plant.

Nozzles should be placed so as to avoid spraying the cotton foliage. A second application may be required during humid growing conditions to control perennials, such as Johnsongrass and Nutsedge. Apply the second application 1 to 3 weeks later. Keep spray off cotton foliage.

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

APPLY ONLY WHEN COTTON IS 3 INCHES HIGH TO BLOOM. DO NOT APPLY AFTER FIRST BLOOM. DO NOT FEED TREATED FOLIAGE TO LIVESTOCK, OR GRAZE TREATED AREAS.

NON-CROP: MICRO MONO Herbicide is useful for controlling Johnsongrass, Nutsedge, Barnyardgrass, and similar weeds on drainage ditch banks, right-of-way, storage yards, and similar non-crop areas.

Mix MICRO MONO Herbicide at a rate of 3 to 6 pints, plus 1 to 2 quarts of a suitable surfactant in 100 gallons of water. Spray the unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, re-apply as required.

This product is toxic to wildlife. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

LAWN AND ORNAMENTAL TURF: MICRO MONO Herbicide is useful for control of Dallisgrass, Sandbur, Bahiagrass, Nutsedge, Barnyardgrass, Chickweed and Wood Sorrel with little or no injury to good lawn grasses. On new lawns, do not treat until after three mowings. Good grasses may be temporarily discolored. Zoysia, Bluegrasses, and Bermuda are quite tolerant. Do not use on St. Augustine or Centipede.

Mow lawns 1 to 1 1/2 inches high before treatment. Mix 1 fluid ounce (2 tablespoons) of MICRO MONO Herbicide and 1/2 fluid ounce (1 tablespoon) of a suitable surfactant in 5 gallons of water and apply to an area of 1,000 square feet. Spray thoroughly to wet all undesirable plants. Repeat applications, 10 to 14 days apart, may be needed for good control.