

DIRECTIONS FOR USE

COTTON: MICRO BANDEX Herbicide is useful as a directed application in cotton for post-emergent control of weeds such as Johnsongrass, Nutsedge, Dallisgrass, Goosegrass, Barnyardgrass, Cocklebur, Sandbur, Puncturevine and Morning Glory. It should be applied at the rate of 2 quarts per acre.

Mix 2 quarts of Micro BANDEX Herbicide in 40 gallons of water and apply as directed spray at 40 gallons per acre. Weed control is most effective at temperatures above 70 degrees F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre. Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage. A second application may be required during humid growing conditions to control perennials such as Johnsongrass and Nutsedge. Apply the second application 1 to 3 weeks later.

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

APPLY ONLY WHEN COTTON IS 6 INCHES HIGH TO FIRST BLOOM. DO NOT APPLY AFTER FIRST BLOOM. DO NOT FEED TREATED FOLIAGE TO LIVESTOCK OR GRAZE TREATED AREAS.

NON-CROP: Micro BANDEX Herbicide is useful for control of Johnsongrass, Nutsedge, Barnyardgrass, Morning Glory and other weeds on drainage ditchbanks, rights-of-way, fence rows, storage yards and other non-crop areas.

Mix 5 quarts of Micro BANDEX Herbicide in 100 gallons of water. Spray unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, reapply as required.

DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION, OR BY ANIMALS, WILDLIFE OR AQUATIC LIFE, OR FOR IRRIGATION PURPOSES.

Net Contents 1 gallon

E.P.A. Reg. No. 4841-54

MICRO BANDEX HERBICIDE

MSMA LIQUID PLUS DIURON
For Selective Post-Emergent Weed Control
Read Entire Label Before Using This Product

ACTIVE INGREDIENTS:

Monosodium Acid Methanearsonate -----	33.9%
Diuron (3-3,4-dichlorophenyl-1, 1-dimethylurea) -----	5.1%
INERT INGREDIENTS: -----	61.0%

Total arsenic (as elemental) all in water soluble form - 15.7%.

Product contains 4 lbs. Monosodium Acid Methanearsonate (7.27 lbs. DSMA Equivalent) and 0.6 lbs. Diuron per gallon plus surfactant.

Caution

Keep out of reach of Children

See side panel for additional cautions.

Manufactured by

Mirc Chemical Company, Inc.

Winnboro, Louisiana 71295

CAUTION

CAUTION: KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near food or feed products. Keep children and domestic animals off treated areas until this material has been washed into the soil. May cause irritation of eyes, nose, throat and skin.

ANTIDOTE: If taken internally, induce vomiting and call a physician at once.


NOTICE: Use of this material in a manner or at any time other than in accordance with directions may produce plant injury, excessive residues, or other undesirable results. Micro Chemical Co., Inc. makes no express or implied warranties other than specified on the label.

GENERAL INFORMATION: Micro BANDEX Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. It is a combination of herbicides and surfactant. It is unnecessary to add other surfactant to the spray solution. Best results are obtained on young, actively growing weeds at temperatures above 70 degrees F.

MIXING INSTRUCTIONS: Shake the Micro BANDEX Herbicide container thoroughly before using. Fill the spray equipment reservoir with water and add the required amount of Micro BANDEX Herbicide with agitation. Continue agitation during application. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank.

THE CONTENTS OF THIS CONTAINER MUST BE SHAKEN THOROUGHLY BEFORE EACH USE UNTIL A UNIFORM MIXTURE IS OBTAINED, AS THE CONTENTS SEPARATE ON STANDING.

CONTAINER DISPOSAL: Do not reuse container. Destroy when empty.



ALL INFORMATION SUBJECT TO CHANGE