2 LBS. NET



*SEVIN • 5 • DUST

Controls The Following Insects:

- Mexican Bean Beetle
- Bean Leaf Beetle
- Flea Beetle
- Leafhopper
- Cucumber Beetle
- Potato Beetle

OUTSTANDING AGAINST
MEXICAN BEAN BEETLE FLEA BEETLES
EFFECTIVE LONG LASTING

Active Ingredients	
*Carbaryl (1-naphthyl N-methylcarbamate)	5%
Inert Ingredients	95%
Total	100%

* Trademark of Union Carbide Corp.

CAUTION: Keep out of reach of children

4876-7



*SEVIN 5 DUST

DIRECTIONS FOR USE

Apply thoroughly to vegetable plants, making sure to treat both top and bottom side of leaves. Make first application when insects first appear. Begin early when new growth is starting. Repeat as necessary.

BEANS: To control Mexican bean beetles and bean leaf beetles, apply 20 to 40 pounds of Sevin 5 dust per acre. Do not use within three days of harvest.

TOMATOES: To control tomato fruit worms, flea beetles and leafhoppers, apply 20 to 40 pounds of Sevin 5 dust per acre.

POTATOES: To control leafhoppers, Colorado potato beetles and potato leafhoppers, apply 20 to 40 pounds of Sevin 5 dust per acre.

MELONS, CUCUMBERS AND SQUASH: To control melon worms, striped and spotted cucumber beetles and flea beetles, apply 20 pounds of Sevin 5 dust per acre.

CAUTION: Harmful if inhaled, swallowed or absorbed through skin. Do not breathe dust, do not take internally. KEEP OUT OF THE REACH OF CHILDREN AND ANIMALS. For the protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

DO NOT REUSE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

U.S.D.A. Reg. No. 4876-7