[illegible]XAMPLE

[illegible]: [illegible] Livestock Pests

[illegible]

[illegible] (Active Ingredient) [illegible].0%
[illegible] 27.0%
[illegible] 4.0%
[illegible] Total 100.0%

[illegible]

SMALL GRAINS ([illegible]) [illegible] spray thoroughly [illegible] give good coverage. Use the higher dosage [illegible] when vegetation is thick and tall. Do not feed treated [illegible] for slaughter.

CORN, WHEAT, OATS, RYE AND BARLEY; [illegible] quarter acre, when spraying [illegible] good coverage. When spraying wheat, [illegible] with enough water for good coverage. [illegible] animals being finished for slaughter.

[illegible] FIELD PASTURE; Do not [illegible] and adjoining areas at rates of [illegible] as grazing pens and arrowrows [illegible] dairy animals or animals being finished [illegible] Birds feeding on treated areas [illegible].

LIVESTOCK SPRAY: [illegible] thoroughly using 1 quart per 100 gallons of water. [illegible] animals being finished for slaughter within [illegible].

STABLE TYPE BACK [illegible]: [illegible] 3 gallons of light oil. Do not use [illegible] treated within 28 days of slaughter.

CAUTION: Do not breathe spray or dust. [illegible] eyes, on skin, or on clothing. Do not store near [illegible] soap and water after handling and before eating [illegible] daily. Do not apply or allow to drift into areas occupied by [illegible] beneficial animals.
This product is toxic to fish and wild life. Keep out of rivers, [illegible] Apply this product only as specified on this label. Rinse empty [illegible] dispose of waste by burying in non-crop lands away from water supplies. [illegible] ainers by punching holes in them and burying with waste [illegible].

[illegible] SUPPLY [illegible]