

USDA Reg. No

**LINDANE  
SPRAY CONCENTRATE**

Net Content -----

**CAUTION:** Keep out of the reach of children  
See right side of panel for additional cautions

Camerasman's Note  
Poor Copy

Active Ingredient		
Lindane (gamma isomer of benzene hexachloride)	10.0%	
Inert Ingredient	90.0%	
	total	100.0%

AG. SUPPLY COMPANY  
Hopkinsville, Ky.

Directions for use on animals

BEEF CATTLE, SHEEP, GOATS, HOGS, AND HORSES-- To control lice, ticks, fleas, and Sarcoptic mange mites, flies and mosquitoes, spray thoroughly or dip with a dilution of one (1) tablespoon per gallon of water (1 pint per 30 gallon of water). Repeat as necessary. Do not use on dairy animals. Do not treat animals under 3 months of age. Do not treat cattle, hogs or sheep within 30 days of slaughter. Do not dip within 60 days of slaughter. Do not use on poultry or in poultry houses.

FARM BUILDING RESIDUAL SPRAY: To control flies, roaches, ants, and mosquitoes use six tablespoons per gallon of water, spray barns, dog kennels, hog pens etc. Spray thoroughly throughout building. Do not spray this dilution on animals. Do not contaminate feed or foodstuffs or drinking water. Exclude animals when spraying. Do not contaminate milk or milking equipment. Do not use in poultry houses.

GRAIN AND OTHER STORAGE BINS FOR STORED GRAINS: To control saw-toothed grain beetle, granary weevil, confused flour beetle, corn or rice weevil, agaveis grain moth and indian meal moth, use a dilution of 1 gallon per 25 gallon of water, applying 2 gallon of diluted spray per 1000 square feet of bin area. Treat empty bins prior to filling.

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with or absorbed through skin. In case of skin contact, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not use on household pets. Do not use, pour or spill near heat or open flame. This product is toxic to fish and wild life. Keep out of rivers, streams, ponds. Use equipment and containers to dispose of waste by burning. Do not use spray or pour on crops. Decontaminate containers by washing with soap and water. Do not use near children. (keep out of smoke).

<b>ACCEPTED</b>
<b>OCT 20 1967</b>
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND REPLICIDE ACT FOR FILING IN CONNECTION WITH ED UNDER NO. 48,76-24 SUBJECT TO ATTACHED COMMENTS.