

Middle Panel

USDA REG. NO. 4876-

Net 1 Pint

SPRAY-AWAY WEED & GRASS KILLER

TOP KILLS WEEDS AND GRASS IN:

Driveways - Walks - Fence Lines, around

Buildings, Trees, Shrubs and Evergreens

Excellent for chemical edging of driveways, walks,
rose beds, and flower beds.

CAUTION: Keep out of the reach of children

See side panel for additional cautions.

ACTIVE INGREDIENTS:

Sodium Cacodylate 12.6%

Dimethylarsinic Acid (Cacodylic Acid) 2.2%


INERT INGREDIENTS: 85.2%

Total Arsenic (as elemental
all in water soluble form) 7.1%

Product contains 1.24 lbs. Cacodylic acid
equivalent per gallon.

"AG" SUPPLY COMPANY
HOPKINSVILLE, KENTUCKY

Front Panel 3" x 4"



Left Panel

GENERAL INFORMATION

useful for general post-emergent weed control. It is non-selective in its action. Its phytotoxic properties are quickly inactivated on contact with soil.


After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.
CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent and discard in a safe place.

DIRECTIONS FOR USE

DRIVEWAYS, WALKS, FENCE LINES: To top kill all vegetation, mix 1 pint of **SPRAY-AWAY** in 3 gallons of water. Apply spray to wet all foliage thoroughly. On perennials, repeat application as necessary.

CLEAN-UP AROUND TREES, SHRUBS, AND EVERGREENS: Mix 1 pint **SPRAY-AWAY** to 3 gallons of water. Direct spray onto vegetation to be eliminated at base of trees and shrubs. **DO NOT APPLY** to or allow spray to contact foliage, green bark, graft unions, or broken bark when spraying around trees or shrubs.

EDGING: For a weed free border along walks, driveways, flower beds, and gardens-- mix 1 pint **SPRAY-AWAY** in 3 gallons of water. Spray to wet all plants in strips you wish to be free of vegetation. **DO NOT ALLOW** spray to contact desired plants.



Right Panel

DIRECTIONS FOR USE: LAWN RENOVATION:
SPRAY-AWAY may be used to kill-out an area to be re-seeded. Mix 1 pint of SPRAY-AWAY in 3 gallons of water and apply to an area of 800 square feet. Completely wet all vegetation to be killed. Bermuda grass and other deep-rooted plants may require a second treatment. Re-apply in five days for such areas.


About 5 days after last treatment, rake away killed foliage. Water well and seed at rate for new lawns. Rake seed into soil, mulch lightly with straw, and keep area well watered.

Keep children and pets off treated areas until after first rain or sprinkling following treatment. Do not track from treated to untreated areas.

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into soil.

ANTIDOTE: If taken internally induce vomiting and call a physician at once.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT



SPRAY-AWAY WEED & GRASS KILLER

KILLS WEEDS AND GRASS IN:

Driveways - Walks - Fence Lines, around

Buildings, Trees, Shrubs and Evergreens

Excellent for chemical edging of driveways, walks,
rose beds, and flower beds.

CAUTION: Keep out of the reach of children
See side panel for additional cautions.

ACTIVE INGREDIENTS:

Sodium Cacodylate 12.6%
Dimethylarsinic Acid (Cacodylic Acid) 2.2%


INERT INGREDIENTS: 85.2%

Total Arsenic (as elemental
all in water soluble form) 7.1%

Product contains 1.24 lbs. Cacodylic acid
equivalent per gallon.

"AG" SUPPLY COMPANY
HOPKINSVILLE, KENTUCKY

Front Panel 5' x 4'



Left Panel

GENERAL INFORMATION

useful for general post-emergent weed control. It is non-selective in its action. Its phytotoxic properties are quickly inactivated on contact with soil.

After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent and discard in a safe place.

DIRECTIONS FOR USE

DRIVEWAYS, WALKS, FENCE LINES: To top kill all vegetation, mix 1 pint of **SPRAY-AWAY** in 3 gallons of water. Apply spray to wet all foliage thoroughly. On perennials, repeat application as necessary.

CLEAN-UP AROUND TREES, SHRUBS, AND EVERGREENS: Mix 1 pint **SPRAY-AWAY** to 3 gallons of water. Direct spray onto vegetation to be eliminated at base of trees and shrubs. DO NOT APPLY to or allow spray to contact foliage, green bark, graft unions, or broken bark when spraying around trees or shrubs.

EDGING: For a weed free border along walks, driveways, flower beds, and gardens-- mix 1 pint **SPRAY-AWAY** in 3 gallons of water. Spray to wet all plants in strips you wish to be free of vegetation. DO NOT ALLOW spray to contact desired plants.

3/1/61

Right Panel

DIRECTIONS FOR USE: LAWN RENOVATION:

SPRAY-AWAY may be used to kill-out an area to be re-seeded. Mix 1 pint of SPRAY-AWAY in 3 gallons of water and apply to an area of 800 square feet. Completely wet all vegetation to be killed. Bermuda grass and other deep-rooted plants may require a second treatment. Re-apply in five days for such areas.

About 5 days after last treatment, rake away killed foliage. Water well and seed at rate for new lawns. Rake seed into soil, mulch lightly with straw, and keep area well watered.

Keep children and pets off treated areas until after first rain or sprinkling following treatment. Do not track from treated to untreated areas.

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into soil.

ANTIDOTE: If taken internally induce vomiting and call a physician at once.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

3/11/68

4/8/68

ACCEPT
JAN 20 / 9
UNDER THE FEDERAL
FUNCTION
FOR ED
ED UNDER
TO ATTACHED

GRO-WELL

GRO-WELL. CRABGRASS AND BROADLEAF WEED KILLER

1.49

KILLS CRABGRASS, DANDELION,
PLANTAIN AND CERTAIN OTHER WEEDS
COVERS UP TO 1,000 SQ. FT.

CAUTION
KEEP OUT OF THE REACH OF CHILDREN
ADDITIONAL CAUTIONS ON REVERSE SIDE

CAUTION:

GRO-WELL CRABGRASS & BROADLEAF WEED KILLER

This will kill crabgrass, chickweed and most Broadleaf Lawn Weeds such as Dandelion, Plantain, Knotweed, Spotted Spurge, Pennywort and Purslane. The best time to apply is in late afternoon and when plants are small and seed heads have not formed. Two or more applications 7 days apart may be needed until plants are dead. Application may be needed again later in season as new seedlings germinate. Read entire label and measure area carefully. The important factor is the amount of chemical applied to a given area. A coarse spray type sprayer that attaches to a hose, such as the GRO-WELL Lawn Sprayer, may be used.

NOTE: 3 Teaspoonfuls = 1 Tablespoonful, 2 Tablespoonfuls = 1 Ounce
FOR TEMPERATURE 80° AND ABOVE: Dilute 8 oz. of chemical with 5 Gallons of water and apply to 1000 sq. ft. of lawn.

FOR TEMPERATURE BELOW 80°: Dilute 16 oz. of chemical with 5 Gallons of water and apply to 1000sq. ft. of lawn.

CAUTION: Soil should be moist and grass in good growing condition before and three days after treatment. If necessary water well day before, skip day of and day after treatment then water next two days. Do not use on newly seeded lawns until after 3rd mowing. Reseeding may be done 3 weeks later. As Bents, Fescues and Clover may be temporarily discolored use with care. Do not use same sprayer for other than lawn spraying. Do not use on St. Augustine, Carpet, or Centipede Grasses or on Bent Golf Greens.

WARNING: Keep out of reach of children and pets and keep them off treated areas. In case of contact with eyes (Flush Profusely with water) or if taken internally (induce Vomiting) contact Doctor immediately. Avoid breathing spray mist and contact with skin or clothing. Wash off immediately with warm soap and water. Avoid contamination of feed and foodstuffs. Do not spray or allow to drift on to desirable Flowers, Trees, Shrubs, Vegetables or into lakes, streams or ponds. Coarse sprays are less likely to drift. Do not store near Fertilizers, Seeds, Insecticides or Fungicides.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

ACTIVE INGREDIENTS	PERCENTAGE
Dodecyl Ammonium Methyl Arsonate	8.0%
Octyl Ammonium Methyl Arsonate	8.00%
Octylamine Salt of 2,4-Dichlorophenoxyacetic Acid	5.44%
INERT INGREDIENTS	78.56%

*Equiv. to 2,4-Dichlorophenoxyacetic Acid 3.43%

Sold by: J. & L. ADIKES, INC., JAMAICA, NEW YORK 11423
NET CONTENTS 8 OUNCES
EPA Reg. No. 5535-25

ACTIVE INGREDIENTS
DODECYLAMMONIUM
OCTYLAMMONIUM METH
* OCTYLAMINE SALT OF 2,
INERT INGREDIENTS.

~~EXPRESSED AS ELEMENTAL~~
TOTAL ARSENIC EXPRESSED AS ELEMENTAL, ALL IN WATER SOLUBLE FORM..... 4.0