

DIRECTIONS FOR USE

IMPORTANT: Read these entire directions before using Tobacco Plant Bed Systemic Granules.

TOBACCO PLANT BEDS: For the control of aphids and flea beetles, apply 1 ounce per 100 square feet of bed. This container will treat 1000 square feet of plant bed.

Apply evenly over bed area, just before seeding. Work the granules into the top 2 or 3 inches of soil and seed beds in normal manner. An alternative treatment may be made by broadcasting the granules evenly over the plants after emergence and watering thoroughly. Do not apply more than once per season.

DANGER

Poisonous if swallowed, inhaled or absorbed through the skin. Rapidly absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wear protective clothing and cotton or natural rubber gloves. If cotton gloves are used, they must be laundered or discarded after each day's use. Do not wear the same gloves for other work. Rubber gloves should be washed with soap and water after use. Do not contaminate feed or food. Keep out of reach of children. Keep all unprotected persons out of the operating area or vicinity. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Do not reuse container in any manner in connection with food, feed or water. Destroy container to prevent reuse.

Wear mask or respirator of a type found by the U.S. Bureau of Mines to give adequate protection against this material.

This product is poisonous to fish and wildlife. Exposed granules on surface of ground may be hazardous to birds. Keep out of lakes, ponds, and streams. Apply this product only as specified on this label.

Notice of Warranty:

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

TREATMENT: In case of poisoning, call a physician immediately. Have patient lie down and keep quiet. If swallowed, induce vomiting by giving a tablespoonful of salt in a glass of warm water. Administer milk or water freely and again induce vomiting until vomit fluid is clear. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

POISON

TO PHYSICIAN:

ANTIDOTE — administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine. Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor-nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

USDA Reg. No. 4876-33

Tobacco Plant Bed

Contains * DI-SYSTON

SYSTEMIC GRANULES
For Control of
APHIDS & FLEA BEETLES



**WORKS INSIDE THE
PLANTS AS THEY GROW**

**DANGER
POISON**



RESTRICTIONS - FOR COMMERCIAL GROWERS ONLY

Not for home garden use. Do not use on crops used for food or storage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse. Not for use or storage in or around the home.

Keep Out Of Reach Of Children
See rear panel for antidote and warning statements.

ACTIVE INGREDIENT

0,0-Diethyl S-(2-(ethylthio)ethyl)
phosphorodithioate† 15%
INERT INGREDIENT 85%

100%

DI-SYSTON is a Reg. TM in U.S. and Canada Pat. Offs. by
Farbenfabriken Bayer, A.G., Chemagro Corporation Licensee.
†U.S. Patent 2,759,010

RE-PACKAGED BY

"AG" SUPPLY COMPANY

HOPKINSVILLE KY.

ACCEPTED WITH COMMENTS

8-25-71
4876-33