

U.S.D.A. Reg. No. 4876-38

**DIRECTIONS FOR USE:**

**IMPORTANT.** The directions on this label are based on tests and field experience relating to (a) effectiveness; and (b) possible injury to plants and animals; and (c) residues in food, feed, meat and milk. Follow directions for most effective and safest use.

**TOBACCO INSECT CONTROL**

**IN FIELDS**—For budworms, flea beetles, hornworms, grasshoppers, use 1/2 to 2 pounds in 25 to 30 gallons of water per acre. Full coverage of plants is essential. Use lower rate on young plants up to knee high; higher rate on mature plants. Begin treatments when worms are small. Avoid excessive application in plant bud as injury may result.

Treated fields may be entered immediately after foliage has dried. If late applications are necessary, allow 3 days before priming or cutting.

**IN PLANT BEDS**—For tobacco flea beetle, use 2.5 level tablespoonfuls per gallon water, or 1/4 pound per 50 gallon water, and apply 6 gallon per 100 square yards. Avoid excessive applications; plant injury may result.

**CAUTION**

**HARMFUL IF INHALED OR SWALLOWED.** Avoid Breathing of Dust or Spray. Do Not Take Internally. Avoid Prolonged or Repeated Contact with Skin.

Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating.

Wash thoroughly after handling.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

**AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.**

**GENERAL CAUTIONS**

1. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

**COMPATIBILITY**

Unstable under highly alkaline conditions. Not effective if used with alkaline materials such as Bordeaux, lime, lime sulfur and casein-lime spreaders. Seller makes no warranty, expressed, or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**"AG" SUPPLY COMPANY**  
HOPKINSVILLE KENTUCKY

NET CONTENTS 2 LBS.



**SEVIN\* 85W**  
**TOBACCO SPRAY**  
**Controls:**



- ★ Flea Beetles
- ★ Hornworms
- ★ Budworms
- ★ Grasshoppers

**CAUTION: Keep Out of Reach of Children**  
See Side Panel For Additional Cautions

**ACTIVE INGREDIENTS:**

Carbaryl-(1-Naphthyl Methylcarbamate)	.85.00%
INERT INGREDIENTS	15.00%
	<hr/>
	100.00%

\*SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide.

Packaged by

**"AG" SUPPLY COMPANY**

**ACCEPTED**

MAY 27 1971