


## DIRECTIONS FOR USE

Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe 5 day interval between last application and harvest.

**TOBACCO:** To control cabbage loopers, budworm, flea beetle, green peach aphid, and hornworm, in fields, apply 1 2 quarts per acre in sufficient water to give thorough coverage, using not less than 15 gallons of water per acre. To control the above mentioned insects in seed bed areas, mix 1 pint in 50 gallons of water applying six gallons of mixture per 100 square yards of plant beds. To control green june bug larvae in plant beds, mix ½ pint in 50 gallons of water, apply 1 gallon of mixture to 1 square yard of plant bed.

**WARNING—**Hazardous if swallowed, inhaled, or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator approved by the U. S. Bureau of Mines for endosulfan protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

# THIODAN TOBACCO SPRAY



**WARNING: Keep Out of the Reach of Children**

SEE SIDE PANEL FOR ADDITIONAL WARNINGS

Active Ingredient

• Endosulfan (Hexachlorohexahydromethano - 2, 4, 3 - benzodioxathiepin oxide) ..... 24.00%

Inert Ingredients ..... 76.00%

Contains 2 pounds of endosulfan per gallon.

\*Thiodan U. S. Patent No. 2,799,685.

Thiodan is a registered trademark of Canadian Roechst Limited.

**WARNING:** Workers entering treated areas within 24 hours of application should wear protective clothing.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.


**DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.** Do not store at temperatures below 20 degrees F. Do not use in uniluted form.

Not for use or storage in or around the home.

Rinse equipment and containers and dispose of waste by burying at least 18" deep in non-crop areas away from water supplies. Do not reuse containers; smash or perforate and bury with waste.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assures all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



USDA  
Reg. No. 4876-39


Net Contents  
1 Gallon

**THIODAN**  
TOBACCO PROTECTANT

Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe 5 day interval between last application and harvest.

**TOBACCO:** To control cabbage loopers, budworm, flea beetle, green peach aphid, and hornworm, in fields, apply 1 to 2 quarts per acre in sufficient water to give thorough coverage, using not less than 15 gallons of water per acre. To control the above mentioned insects in seed bed areas, mix 1 pint in 50 gallons of water applying six gallons of mixture per 100 square yards of plant beds. To control green June bug larvae in plant beds, mix 1/2 pint in 50 gallons of water, apply 1 gallon of mixture to 1 square yard of plant bed.

**WARNING**—Hazardous if swallowed, inhaled, or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator approved by the U. S. Bureau of Mines for endosulfan protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.



**SEE SIDE PANEL FOR ADDITIONAL WARNINGS**

**Active Ingredient**

• Endosulfan Hexachlorocyclohexylhydromethano — 2, 4, 3 —  
benzodioxathiopyridine oxide ..... 24.90%

Inert Ingredients ..... 75.10%

Contains 2 pounds of endosulfan per gallon.

• Thiodan U. S. Patent No. 2,799,685.

Thiodan is a registered trademark of Canadian Rochem Limited.

Packaged By

**"AG" SUPPLY CO.**  
HOPKINSVILLE, KY. 42240

**WARNING:** Workers entering treated areas within 24 hours of application should wear protective clothing.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

**DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.** Do not store at temperature below 20 degrees F. Do not use in undiluted form.

Not for use or storage in or around the home.

Rinse equipment and containers and dispose of waste by burying at least 18" deep in non-crop areas away from water supplies. Do not reuse containers; smash or perforate and bury with waste.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions. To cause animal, plant or crop injury, illegal residues or other undesirable results may occur.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.