



4 LBS. NET




CONTAINS: \* SEVIN • ROTENONE • ZINEB

CONTROLS INSECTS

CONTROLS DISEASE



**INSECTS**  
 BEAN LEAF BEETLE  
 MEXICAN BEAN BEETLE  
 CABBAGE WORM  
 CUCUMBER BEETLE  
 FIF A BEETLE  
 APHIDS (PLANT LICE)



**DISEASES**  
 TOMATO BLIGHTS  
 ANTHRACNOSE  
 LEAFSPOTS  
 BEAN RUST  
 DOWNY MILDEW

**ACTIVE INGREDIENTS**

* Carbaryl (1-Naphthyl N-Methylcarbamate)	2.00 ✓
Rotenone	1.00 ✓
Other cube resins	2.00 ✓
Zineb (Zinc Ethylene Bisdithiocarbamate)	5.00 ✓

**INERT INGREDIENTS**

	90.00 ✓
Total	100.00 ✓

\* Trademark of Union Carbide Corp.

**CAUTION:** Keep out of reach of children.  
 SEE REAR PANEL FOR ADDITIONAL CAUTION

**TRIPLE ACTION VEGETABLE DUST**  
 DIRECTIONS FOR USE

**DUST:** Apply thoroughly to vegetables and flowering plants, making sure to treat both top and bottom sides of leaves, stems and fruit. Make first application when insects first appear and / or disease is anticipated. Begin early, when new growth is starting. Repeat as necessary, generally at 7 day intervals.

**BEANS:** (Snap and Lima)—Controls leafhoppers, Mexican bean beetles, bean leaf beetles and thrips. Diseases: downy mildew, bean rust, and anthracnose. Do not apply within 7 days of harvest.

**CABBAGE, CAULIFLOWER, BROCCOLI, TURNIPS AND CARROTS**—Controls imported cabbage worm, cabbage looper, flea beetles. Diseases: downy mildew and alternaria leaf spot. Stop use 14 days before harvest on Broccoli, 7 days on Cabbage and Cauliflower, 7 days on Turnips and Carrots, or 14 days if tops are to be used for feed or food.

**MELONS, CUCUMBERS AND SQUASH**—Controls melon worms, striped and spotted cucumber beetles and flea beetles. Diseases: downy mildew, alternaria (Macrosporium blight), anthracnose and gummy stem blight. Stop use 7 days before harvest.

**TOMATOES AND EGGPLANT**—Controls tomato horn worm, tomato fruit worm, flea beetles, leafhoppers. Diseases include early and late blight, anthracnose, stemphylium and septoria leaf spots and cladosporium gray mold. Stop use 7 days before harvest.

**FLOWERS AND ORNAMENTALS** (On roses, mums, snapdragons, glads, hollyhocks) —Controls botrytis blight and stemphylium on glads, rust, septoria leaf spot and anthracnose on snaps, mums, carnations and hollyhocks. Black spot on roses. Kills thrips, grasshoppers and cutworms. For cutworms, dust the soil around the base of each plant.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

**CAUTION:** HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid contact. Wash hands before eating.

**DO NOT REUSE EMPTY CONTAINER.** Destroy it by burying with waste or burning. Stay away from smoke or fumes.

12-29-71  
 4876-42