

E.P.A. Reg. No. 4876-47

Net Content 2 Oz.



STREPTOMYCIN SULFATE

For control of
Blue Mold & Wildfire on Tobacco
Fire Blight on Apples & Pears

Active Ingredient (streptomycin sulfate).....21.2%
Inert Ingredient.....78.8%

For Agricultural Use Only

ACCEPTED

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR RECORD
REGISTERED UNDER NO. 4876-47

Apply this fungicide to plants as directed below, and apply with a fine spray to the growing parts of the plants.

Application Rate Water Application

When plants are the size of a dime or when blight is first reported in the area.

Blue Mold

Spray at 5-7 day intervals until plants are set into the field until the blight is no longer a problem.

Spray trees every 3-4 days during blossom time. After petal fall, apply spray every 10-14 days to control wild blight. Do not apply within 30 days of harvest or spray 10-15 days of petal fall on apples.

Do not use this fungicide on plants that are known to be susceptible to streptomycin resistance. Do not use this fungicide on plants that are known to be susceptible to streptomycin resistance. Do not use this fungicide on plants that are known to be susceptible to streptomycin resistance.

Do not use this fungicide on plants that are known to be susceptible to streptomycin resistance.