

DIRECTIONS FOR USE

| CROP | INSECT | Amount of Chlordane per acre | |
|---|---|------------------------------|----------------|
| | | Row | Broadcast |
| CORN (Field & Sweet) | Corn Rootworm (Northern and Southern) | 1 pt. | 2 pts. |
| | Corn Field Ants Corn Root Aphids | | |
| CORN (Field & Sweet) PEANUTS TOBACCO | Seed Corn Beetle Larvae Seed Corn Maggot Wireworms White Grubs | 2 pts. | 2 1/2 - 4 pts. |
| | Cutworms | | 2 1/2 - 4 pts. |
| | Japanese Beetle Larvae | | 5 - 10 pts. |
| | Asiatic Garden Beetle Larvae | | 2 1/2 - 5 pts. |
| | Green June Beetle Larvae | | 3 - 5 pts. |
| | White-fringed Beetle Larvae | | 2 1/2 - 5 pts. |
| | Cutworms | | 2 1/2 - 4 pts. |
| VEGETABLE CROPS Beans Beets Blackeyed Peas Broccoli Brussel Sprouts Cabbage Cantaloupe Cauliflower Celery Collards Cowpeas Cucumber Eggplant Kale Kohlrabi Lettuce Melons Okra Onions Peas Peppers Pimentos Potatoes Radishes Rutabagas Sweet Potatoes Turnips Watermelons | Wireworms White Grubs | 2 pts. | 2 1/2 - 4 pts. |
| | Asiatic Garden Beetle Larvae White-fringed Beetle Larvae | | 2 1/2 - 5 pts. |
| | Japanese Beetle Larvae | | 5 - 10 pts. |
| | Flea Beetle Larvae (potatoes and sweet potatoes) | | 2 1/2 - 5 pts. |

IMPORTANT

On field and vegetable crops make preplant broadcast or row (where applicable) application to soil after plowing and disc or incorporate into soil immediately.

Do not graze small grain grown on treated soil.

Where a range of rates is given, use the higher rate of Chlordane on muck or highly organic soil.



Net Contents 1 Gallon


CHLORDANE 72EC SOIL INSECTICIDE

ACTIVE INGREDIENTS:

Technical Chlordane 72%
Petroleum Distillate 21%

INERT INGREDIENTS 7%
TOTAL 100%

Equivalent to 43.2% Octachloro-4,7-methanotetrahydroindane and 28.8% related compounds.



WARNING

KEEP OUT OF REACH OF CHILDREN

See label panel for additional information and directions for use.

E.F.A. NO. 4876-50


WARNING

May be fatal if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Keep children and pets off treated turf until material has been washed into the soil by watering or rain.

Toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

ANTIDOTE




| | | |
|--|--|---|
| Black-eyed Peas, Broccoli, Brussel Sprouts, Cabbage, Cantaloupe, Cauliflower, Celery, Collards, Cowpeas, Cucumber, Eggplant, Kale, Kohlrabi, Lettuce, Melons, Okra, Onions, Peas, Peppers, Pimientos, Potatoes, Radishes, Rutabagas, Sweet Potatoes, Turnips, Watermelons | White Grubs Asiatic Garden Beetle Larvae White-fringed Beetle Larvae Japanese Beetle Larvae Flea Beetle Larvae (potatoes and sweet potatoes) | 2 pts. 2 1/2 - 4 pts. 5 - 10 pts. 2 1/2 - 5 pts. |
|--|--|---|

IMPORTANT

For field and vegetable crops make preplant broadcast or row (where applicable) application 1-3" after plowing and disc or incorporate into soil immediately.

Do not graze small grain grown on treated soil.

Where a range of rates is given, use the higher rate of Chlordane on muck or highly organic soil.



Equivalent to 43.2% Octachloro-4,7-
methanotetrahydroindane and
28.8% related compounds.

WARNING

KEEP OUT OF REACH OF CHILDREN

See label for directions, cautions, and first aid.

E.P.A. No. 4876-50

SEE THE BOTTOM

WARNING

May be fatal if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Keep children and pets off treated turf until material has been washed into the soil by watering or rain.

Toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

ANTIDOTE

EXTERNAL---In case of spillage on skin, wash promptly with soap and water.

INTERNAL---Emetic of 1 tablespoon mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL

Do not reuse empty container. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. Keep out of sight.

NOT-WARRANTY: Seller makes no warranty, expressed or implied, regarding this product except the composition as set forth in the ingredient statement herein. Buyer and user assume all risk of possession, handling or use whether or not in accordance with any directions contained herein.

