

GENERAL INFORMATION

Diazinon AG500 TOBACCO INSECTICIDE should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil, depending on the type of application.

DIRECTIONS FOR USE


TOBACCO: APHIDS, FLEA BEETLES: Apply 1/2-3/4 pt. per acre. Do not apply within 3 days of harvest.

CUTWORMS: Broadcast 2-3 qts. per acre before transplanting and work into the soil 2-4 inches immediately.

GREEN JUNE BEETLE LARVAE: Mix 1/4 pt. in 50 gals. of water and apply as a drench (100 gals. per 100 sq. yds.) to uprooted areas of plant beds with sprinkling can.

WIREWORMS: Broadcast 2-3 qts. per acre, 3-4 weeks before transplanting. Work into soil 6-9 inches immediately.

GRASSHOPPERS: Apply 1 pt. per acre. Do not apply within 14 days of harvest.



WARNING: May be fatal if swallowed. May be absorbed through the skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Do not permit children or pets to go on to sprayed grass until spray has completely dried.

FISH AND WILDLIFE CAUTION: This product is toxic to fish, birds and other wildlife. Eirds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

Do not use, pour, spill or store near heat or open flame.

Seller makes no warranty, expressed or implied concerning the use of this product other than indicated or the label. Buyer assumes all risk of use and/or landling of this material when such use and/or handling is contrary to label instructions.

2-3 qts. per acre bend work into the soil ly.

WAE: Mix 1/4 pt. in ad apply as a drench eq. yds.) to uprooted with sprinkling can.

t 2-3 qts. per acre, nsplanting. Work into diately.

l pt. per acre. Do days of harvest.

-FOR CONTROL OF APHIDS • FLEA BEETLES • WIREWORMS CUTWORMS and GRASSHOPPERS ON TOBACCO

MAY BE ADDED TO OTHER TOBACCO INSECTICIDES FOR IMPROVED CONTROL OF FLEA BEETLES AND APHIDS.

Active Ingredients:

*Diazinon * — registered trademark of CIBA-GEIGY Corporation.

*O,O-diethyl O-(2-isopropyl-6-methyl-	
4-pyrimidinyl) phosphorothioate	48 %
Aromatic petroleum derivative solvent	
Inert Ingredients:	
	100%

WARNING: Keep out of reach of Children
See side panel for additional cautions

MANUFACTURED BY

"AG" SUPPLY CO.

HOPKINSVILLE, KENTUCKY





cleaning of equipment o Apply this product only label.

This product is highly to direct treatment or residence tive information may be Cooperative Agricultural

Do not reuse empty contaperforating or crushing a safe place.

Do not use, pour, spil or open flame.

Seller makes no warranty concerning the use of the indicated or the label. But use and/or handling of this and/or handling is contrained.