

Net Wt. 1 Lb.

### DIRECTIONS

Make application of this dust when insects first appear. Dust the plants thoroughly applying to undersides as well as tops of surfaces of foliage and repeat application every 4 to 5 days when necessary. Apply at rate of 30 to 40 lbs. per acre on vegetables and small fruit.

Use the above directions to control:

**APHIDS** and **SPIDER MITES** on TOMATOES, PEAS, BEANS, CABBAGE, BROCCOLI and related cole crops, EGGPLANT, CUCUMBER, SQUASH, MELON, POTATOES, STRAWBERRIES, LETTUCE, SPINACH, TOBACCO, ORNAMENTALS.

**THRIPS** on ONIONS

**MEXICAN BEAN BEETLE, LEAFHOPPER** on BEANS

**IMPORTED CABBAGE WORM** and **CABBAGE LOOPER** ON CABBAGE, BROCCOLI and related cole crops.

**BIRCH LEAF MINER, FOUR LINED LEAF BUGS, LEAFHOPPERS, MEALY BUGS, TARNISHED PLANT BUGS, THRIPS, WHITE FLY, EUROPEAN PINE SHOOT MOTH, JAPANESE BEETLE (ADULTS), BAGWORMS** on ORNAMENTALS.


**WARNING:** Do not apply to head lettuce within 7 days of harvest. Do not apply to Spinach, Tobacco, Cabbage, and related cole crops within 7 days of harvest. Do not apply to leaf lettuce within 14 days of harvest. Do not apply to Cucurbits unless plants are dry. Do not apply to other food crops within 3 days of harvest. Avoid use of the product on certain ferns, including Boston, Maidenhair and Pteris as well as on some species of Crassula.

**CONTAINER DISPOSAL:** Do not reuse empty containers. Crush and bury at least 18 inches deep in a safe place away from water supplies.

**CAUTION:** THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT OR RESIDUES ON CROPS. PROTECTIVE INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE.

EPA Reg. No. 4887-17

EPA Est. 4887-GA-1



# STEPHENSON

## 4% MALATHION DUST

(PREMIUM GRADE)

These instructions apply to use on vegetables and small fruit.

**CAUTION:** (SEE CAUTION LABEL)

ACTIVE INGREDIENT	.....	4%
*MALATHION	.....	
INERT INGREDIENTS	.....	96%
TOTAL	.....	100%

\*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate

### CAUTION

AVOID CONTACT WITH SKIN  
WASH THOROUGHLY AFTER USE  
DO NOT CONTAMINATE FOOD AND FEEDSTUFFS

Do not use on plants treated with cholinesterase inhibitors. Atropine is antidote.

For limited quantities the use of this product is restricted to certain crops and areas.

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK, GEORGIA

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK, GEORGIA

### DIRECTIONS

Mix 1/2 lb. of this dust with 100 lbs. of water. Spray thoroughly, applying to undersides of leaves as well as top of fruit. Repeat application every 4 to 5 days when necessary. Apply at 100 to 200 lbs. per acre on vegetable crops and small fruit.

Use the above directions to control:

**APHIDS** and **SPIDER MITES** on TOMATOES, PEAS, BEANS, CABBAGE, BROCCOLI and related cole crops, EGGPLANT, CUCUMBER, SQUASH, MELON, POTATOES, STRAWBERRIES, LETTUCE, SPINACH, TOBACCO, ORNAMENTALS.

**THRIPS** on ONIONS

**MEXICAN BEAN BEETLE, LEAFHOPPER** on BEANS

**IMPORTED CABBAGE WORM** and **CABBAGE LOOPER** on CABBAGE, BROCCOLI and related cole crops.

**BIRCH LEAF MINER, FOUR LINED LEAF BUGS, LEAFHOPPERS, MEALY BUGS, TARNISHED PLANT BUGS, THRIPS, WHITE FLY, EUROPEAN PINE SHOOT MOTH, JAPANESE BEETLE (ADULTS), BAGWORMS** on ORNAMENTALS.

**WARNING:** Do not apply to head lettuce within 7 days of harvest. Do not apply to Spinach, Tobacco, Cabbage, and related cole crops within 7 days of harvest. Do not apply to leaf lettuce within 14 days of harvest. Do not apply to Cucurbits unless plants are dry. Do not apply to other food crops within 14 days of harvest. Avoid use of the product on certain ferns, such as Boston fern, and on other ferns as well as on some species of Crassula.

**CONTAINER DISPOSAL:** Do not reuse empty container. Use hole punch at least 18 inches deep in a safe place away from water supplies.

**CAUTION:** THIS PRODUCT IS HIGHLY TOXIC TO BEES. AVOID USE IN DIRECT TREATMENT OR RESIDUES ON CROPS. PROTECTIVE INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE.

EPA Reg. No. 101-101-01


### CAUTION

**HARMFUL IF SWALLOWED** | **AVOID CONTACT WITH SKIN**  
**KEEP OUT OF REACH OF CHILDREN** | **WASH THOROUGHLY AFTER USE**  
**AVOID BREATHING OF DUST** | **DO NOT CONTAMINATE FOOD AND FEEDSTUFFS**

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on label.

**NOTE TO PHYSICIANS:** Malathion may cause cholinesterase inhibition. Atropine is antidotal.

**NOTICE:** Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.



# STEPHENSON

## Chemicals

# 4% MALATHION DUST

### (PREMIUM GRADE)

Those Insects Listed in Directions on Vegetables and Ornamentals

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION: (SEE CAUTIONS ON SIDE PANEL)**

ACTIVE INGREDIENT  
MALATHION  
INERT INGREDIENTS

\*O, O-dimethyl phosphorothioate of diethylmercaptoacetate

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK, GEORGIA

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK, GEORGIA