

#### **GENERAL DIRECTIONS**


Place bait in locations where rats and mice feed, water or travel. Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable. Bait should be placed in locations not accessible to children, pets, wildlife, and domestic animals or in tamper-proof boxes. Where a continuous source of infestation is present (nearby dumps), permanent bait stations should be established and replenished as needed. **NOTE**, Baits that are sour, moldy or otherwise decomposed should be replaced promptly.

#### **RAT CONTROL**

Place 4 oz. to 1 lb. of STEPHENSON CHEMICALS FUMARIN RAT BAIT where rats are known to run or feed or water. Baiting must be continuous until complete lack of feeding is noticed, USUALLY 10 TO 14 DAYS. NEVER LET A BAIT STATION REMAIN EMPTY MORE THAN 48 HOURS.

#### **MOUSE CONTROL**

Use 1 or 2 tablespoons of bait at various 8 to 12 ft. intervals for control of mice. Make sure that bait is available for at least 15 days.



**GENERAL DIRECTIONS**

... .. NOTE

**RAT CONTROL**

... .. Baiting must be

**MOUSE CONTROL**

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**Container Disposal:** Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.  
USDA Reg. No. 4887-60




A 1 lb. quantity  
of this product  
is sufficient to  
control 4 or 5  
rats or several  
mice.



1M 7-77

Net Wt. ....



**STEPHENSON**

**FUMARIN RAT BAIT  
PELLETS**

**CAUTION: KEEP OUT OF REACH OF CHILDREN.  
(SEE CAUTION ON SIDE PANEL)**

ACTIVE INGREDIENT	
3 (alpha-acetyl) furfuryl-4 hydroxycoumarin	0.025%
INERT INGREDIENTS	99.975%
Total	100.000%

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK, GEORGIA

0.1 lb. quantity  
of this product  
is sufficient to  
control 4 or 5  
rats or several  
mice.



**CAUTION**

THE ACTIVE INGREDIENT, FUMARIN, IS AN ANTI-COAGULANT CHEMICAL WHICH MAY CAUSE SERIOUS HEMORRHAGE IN HUMANS, PETS, WILDLIFE OR LIVESTOCK. IF EATEN, GIVE A TABLESPOON OF SALT IN A GLASS OF WARM WATER AND REPEAT UNTIL VOMIT FLUID IS CLEAR. CALL A PHYSICIAN IMMEDIATELY.

**NOTE FOR PHYSICIANS**

In case of ingestion of FUMARIN, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol.

STEPHENSON  
Chemicals

**FUMARIN RAT BAIT  
PELLETS**

**KILLS RATS and MICE**

Cheese Odor Added

**CAUTION:** KEEP OUT OF REACH OF CHILDREN.  
SEE CAUTION ON SIDE PANEL

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK, GEORGIA